

# CONTROLLABILITY FOR SCHRÖDINGER TYPE SYSTEM WITH MIXED DISPERSION ON COMPACT STAR GRAPHS

ROBERTO DE A. CAPISTRANO-FILHO, MÁRCIO CAVALCANTE, AND FERNANDO A. GALLEGÓ

ABSTRACT. In this work we are concerned with solutions to the linear Schrödinger type system with mixed dispersion, the so-called biharmonic Schrödinger equation. Precisely, we are able to prove an exact control property for these solutions with the control in the energy space posed on an oriented star graph structure  $\mathcal{G}$  for  $T > T_{min}$ , with

$$T_{min} = \sqrt{\frac{\bar{L}(L^2 + \pi^2)}{\pi^2\varepsilon(1 - \bar{L}\varepsilon)}},$$

when the couplings and the controls appear only on the Neumann boundary conditions.

## 1. INTRODUCTION

The fourth-order nonlinear Schrödinger (4NLS) equation or biharmonic cubic nonlinear Schrödinger equation

$$(1.1) \quad i\partial_t u + \partial_x^2 u - \partial_x^4 u = \lambda|u|^2 u,$$

have been introduced by Karpman [24] and Karpman and Shagalov [25] to take into account the role of small fourth-order dispersion terms in the propagation of intense laser beams in a bulk medium with Kerr nonlinearity. Equation (1.1) arises in many scientific fields such as quantum mechanics, nonlinear optics and plasma physics, and has been intensively studied with fruitful references (see [9, 24] and references therein).

In the past twenty years such 4NLS have been deeply studied from different mathematical points of view. For example, Fibich *et al.* [21] worked various properties of the equation in the subcritical regime, with part of their analysis relying on very interesting numerical developments. The well-posedness problem and existence of the solutions has been shown (see, for instance, [30]) by means of the energy method, harmonic analysis, etc.

**1.1. Dispersive models on star graphs.** The study of nonlinear dispersive models in a metric graph has attracted a lot of attention of mathematicians, physicists, chemists and engineers, see for details [10, 11, 13, 27] and references therein. In particular, the framework prototype (graph-geometry) for description of these phenomena have been a *star graph*  $\mathcal{G}$ , namely, on metric graphs with  $N$  half-lines of the form  $(0, +\infty)$  connecting at a common vertex  $\nu = 0$ , together with a nonlinear equation suitably defined on the edges such as the nonlinear Schrödinger equation (see Adami *et al.* [1, 2] and Angulo and Goloshchapova [3, 4]). We note that with the introduction of nonlinearities in the dispersive models, the network provides a nice field, where one can look for interesting soliton propagation and nonlinear dynamics in general. A central point that makes this analysis a delicate problem is the presence of a vertex where the underlying one-dimensional star graph should bifurcate (or multi-bifurcate in a general metric graph).

Looking at other nonlinear dispersive systems on graph structure, we have some interesting results. For example, related with well-posedness theory, the second author in [16], studied the local well-posedness for the Cauchy problem associated to Korteweg-de Vries equation in a metric star graph with three semi-infinite edges given by one negative half-line and two positives half-lines attached to a common vertex  $\nu = 0$  (the  $\mathcal{Y}$ -junction framework). Another nonlinear dispersive

---

2020 *Mathematics Subject Classification.* 35R02, 35Q55, 35G30, 93B05, 93B07.

*Key words and phrases.* Exact controllability, Schrödinger type equation, Star graph, Neumann boundary conditions.

equation, the Benjamin–Bona–Mahony (BBM) equation, is treated in [12, 29]. More precisely, Bona and Cascaval [12] obtained local well-posedness in Sobolev space  $H^1$  and Mugnolo and Rault [29] showed the existence of traveling waves for the BBM equation on graphs. Using a different approach Ammari and Crépeau [7] derived results of well-posedness and, also, stabilization for the Benjamin–Bona–Mahony equation in a star-shaped network with bounded edges.

Recently, in [15], the authors deals to present answers for some questions left in [14] concerning the study of the cubic fourth order Schrödinger equation in a star graph structure  $\mathcal{G}$ . Precisely, they considered  $\mathcal{G}$  composed by  $N$  edges parameterized by half-lines  $(0, +\infty)$  attached with a common vertex  $\nu$ . With this structure the manuscript studied the well-posedness of a dispersive model on star graphs with three appropriate vertex conditions.

Regarding to the control theory and inverse problems, let us cite some previous works on star graphs. Ignat *et al.* in [23] worked on the inverse problem for the heat equation and the Schrödinger equation on a tree. Later on, Baudouin and Yamamoto [8] proposed a unified - and simpler - method to study the inverse problem of determining a coefficient. Results of stabilization and boundary controllability for KdV equation on star-shaped graphs was also proved in [6, 17, 18]. Finally, recently, Duca in [19, 20] showed the controllability of the bilinear Schrödinger equation defined on a compact graph. In booth works, with different main goals, the author showed control properties for this system.

We caution that this is only a small sample of the extant work on graphs structure for partial differential equations.

**1.2. Functional framework.** Let us define the graphs  $\mathcal{G}$  given by the central node 0 and edges  $I_j$ , for  $j = 1, 2, \dots, N$ . Thus, for any function  $f : \mathcal{G} \rightarrow \mathbb{C}$ , we set  $f_j = f|_{I_j}$ ,

$$L^2(\mathcal{G}) := \bigoplus_{j=1}^N L^2(I_j) := \{f : \mathcal{G} \rightarrow \mathbb{R} : f_j \in L^2(I_j), j \in \{1, 2, \dots, N\}\}, \quad \|f\|_2 = \left( \sum_{j=1}^N \|f_j\|_{L^2(I_j)}^2 \right)^{1/2}$$

and

$$(f, g)_{L^2(\mathcal{G})} := \operatorname{Re} \int_{-l_1}^0 f_1(x) \overline{g_1(x)} dx + \operatorname{Re} \sum_{j=2}^N \int_0^{l_j} f_j(x) \overline{g_j(x)} dx.$$

Also, we need the following spaces

$$H_0^m(\mathcal{G}) := \bigoplus_{j=1}^N H_0^m(I_j) := \{f : \mathcal{G} \rightarrow \mathbb{C} : f_j \in H^m(I_j), j \in \{1, 2, \dots, N\}\},$$

where  $\partial_x^j f_1(-l_1) = \partial_x^j f_j(l_j) = 0$ ,  $j \in \{1, 2, \dots, m-1\}$  and  $f_1(0) = \alpha_j f_j(0)$ ,  $j \in \{1, 2, \dots, N\}$ , with

$$\|f\|_{H_0^m(\mathcal{G})} = \left( \sum_{j=1}^N \|f_j\|_{H^m(I_j)}^2 \right)^{1/2},$$

for  $m \in \mathbb{N}$  with the natural inner product of  $H_0^s(I_j)$ . We often write,

$$\int_{\mathcal{G}} f dx = \int_{-l_1}^0 f_1(x) dx + \sum_{j=2}^N \int_0^{l_j} f_j(x) dx$$

Then the inner products and the norms of the Hilbert spaces  $L^2(\mathcal{G})$  and  $H_0^m(\mathcal{G})$  are defined by

$$\begin{aligned} \langle f, g \rangle_{L^2(\mathcal{G})} &= \operatorname{Re} \int_{\mathcal{G}} f(x) \overline{g(x)} dx \quad \text{and} \quad \|f\|_{L^2(\mathcal{G})}^2 = \int_{\mathcal{G}} |f|^2 dx, \\ \langle f, g \rangle_{H_0^m(\mathcal{G})} &= \operatorname{Re} \sum_{k \leq m} \int_{\mathcal{G}} \partial_x^k f(x) \overline{\partial_x^k g(x)} dx \quad \text{and} \quad \|f\|_{H_0^m(\mathcal{G})}^2 = \sum_{k \leq m} \int_{\mathcal{G}} |\partial_x^k f|^2 dx. \end{aligned}$$

We will denote  $H^{-s}(\mathcal{G})$  the dual of  $H_0^s(\mathcal{G})$ . By using Poincaré inequality, it follows that

$$\|f\|_{L^2(\mathcal{G})}^2 \leq \frac{L^2}{\pi^2} \|\partial_x f\|_{L^2(\mathcal{G})}^2, \quad \forall f \in H_0^1(\mathcal{G}),$$

where  $L = \max_{j=1,2,\dots,N} \{l_j\}$ . Thus, we have that

$$(1.2) \quad \sum_{k=1}^m \|\partial_x^k f\|_{L^2(\mathcal{G})}^2 \leq \|f\|_{H_0^m(\mathcal{G})}^2 \leq \left( \frac{L^2}{\pi^2} + 1 \right) \sum_{k=1}^m \|\partial_x^k f\|_{L^2(\mathcal{G})}^2.$$

**1.3. Setting of the problem and main result.** Let us now present the problem that we will study in this manuscript. Due the results presented in work [15], naturally, we should see what happens for the control properties for a linear Schrödinger type system with mixed dispersion on compact graph structure  $\mathcal{G}$  of  $(N+1)$  edges  $e_j$  (where  $N \in \mathbb{N}^*$ ), of lengths  $l_j > 0, j \in \{1, \dots, N+1\}$ , connected at one vertex that we assume to be 0 for all the edges. Precisely, we assume that the first edge  $e_1$  is parametrized on the interval  $I_1 := (-l_1, 0)$  and the  $N$  other edges  $e_j$  are parametrized on the interval  $I_j := (0, l_j)$ . On each edge we pose a linear biharmonic NLS equation ( $Bi - NLS$ ). On the first edge ( $j = 1$ ) we put no control and on the other edges ( $j = 2, \dots, N+1$ ) we consider Neumann boundary controls (see Fig. 1).




FIGURE 1. A compact graph with  $N+1$  edges

Thus, in this work, we consider the following system

$$(1.3) \quad \begin{cases} i\partial_t u_j + \partial_x^2 u_j - \partial_x^4 u_j = 0, & (x, t) \in I_j \times (0, T), \quad j = 1, 2, \dots, N \\ u_j(x, 0) = u_{j0}(x), & x \in I_j, \quad j = 1, 2, \dots, N \end{cases}$$

with appropriated boundary conditions as follows

$$(1.4) \quad \begin{cases} u_1(-l_1, t) = \partial_x u_1(-l_1, t) = 0, \\ u_j(l_j, t) = 0, & j \in \{2, 3, \dots, N\}, \\ \partial_x u_j(l_j, t) = h_j(t), & j \in \{2, 3, \dots, N\}, \\ u_1(0, t) = \alpha_j u_j(0, t), & j \in \{2, 3, \dots, N\}, \\ \partial_x u_1(0, t) = \sum_{j=2}^N \frac{\partial_x u_j(0, t)}{\alpha_j} \\ \partial_x^2 u_1(0) = \alpha_j \partial_x^2 u_j(0, t), & j \in \{2, 3, \dots, N\}, \\ \partial_x^3 u_1(0, t) = \sum_{j=2}^N \frac{\partial_x^3 u_j(0, t)}{\alpha_j}. \end{cases}$$

Here  $u_j(x, t)$  is the amplitude of propagation of intense laser beams on the edge  $e_j$  at position  $x \in I_j$  at time  $t$ ,  $h_j = h_j(t)$  is the control on the edge  $e_j$  ( $j \in \{2, \dots, N+1\}$ ) belonging to  $L^2(0, T)$

and  $\alpha_j$  ( $j \in \{2, \dots, N+1\}$ ) is a positive constant. The initial data  $u_{j0}$  are supposed to be  $H^{-2}(\mathcal{G})$  functions of the space variable.

With this framework in hand, our work deals with the following classical control problem.

**Boundary controllability problem:** *For any  $T > 0$ ,  $l_j > 0$ ,  $u_{j0} \in H^{-2}(\mathcal{G})$  and  $u_T \in H^{-2}(\mathcal{G})$ , is it possible to find  $N$  Neumann boundary controls  $h_j \in L^2(0, T)$  such that the solution  $u$  of (1.3)-(1.4) on the tree shaped network of  $N+1$  edges (see Fig. 1) satisfies*

$$(1.5) \quad u(\cdot, 0) = u_0(\cdot) \quad \text{and} \quad u(\cdot, T) = u_T(\cdot)?$$

The answer for that question is given by the following result.

**Theorem 1.1.** *For  $T > 0$  and  $l_1, l_2, \dots, l_N$  positive real numbers, let us suppose that*

$$(1.6) \quad T > \sqrt{\frac{\bar{L}(L^2 + \pi^2)}{\pi^2 \varepsilon (1 - \bar{L} \varepsilon)}} := T_{min}$$

where

$$(1.7) \quad L = \max \{l_1, l_2, \dots, l_N\}, \quad \bar{L} = \max \{2l_1, \max \{l_2, l_3, \dots, l_N\} + l_1\},$$

and

$$(1.8) \quad 0 < \varepsilon < \frac{1}{\bar{L}}.$$

Additionally, suppose that the coefficients of the boundary conditions (1.4) satisfies

$$(1.9) \quad \sum_{j=2}^N \frac{1}{\alpha_j^2} = 1 \quad \text{and} \quad \frac{1}{\alpha_j^2} \leq \frac{1}{N-1}.$$

Then for any  $u_0, u_T \in H^{-2}(\mathcal{G})$ , there exists a control  $h_j(t) \in L^2(0, T)$ , for  $j = 2, \dots, N$ , such that the unique solution  $u(x, t) \in C([0, T]; H^{-2}(\mathcal{G}))$  of (1.3)-(1.4), with  $h_1(t) = 0$ , satisfies (1.5).

**1.4. Outline and structure of the paper.** In this article we prove the exact controllability of the Schrödinger type system with mixed dispersion in star graph structure  $\mathcal{G}$  of  $(N+1)$  edges  $e_j$  of lengths  $l_j > 0$ ,  $j \in \{1, \dots, N+1\}$ , connected at one vertex that we assume to be 0 for all the edges (see Fig. 1). Precisely, we are able to prove that solutions of adjoint system associated to (1.3), with boundary conditions (1.4), preserve conservation laws in  $L^2(\mathcal{G})$ ,  $H^1(\mathcal{G})$  and  $H^2(\mathcal{G})$  (see Appendix A), which are proved *via* Morawetz multipliers<sup>1</sup>. With this in hand, an *observability inequality* associated with the solution of the adjoint system is proved. Here, the relation between  $T > T_{min}$ , where

$$T_{min} = \sqrt{\frac{\bar{L}(L^2 + \pi^2)}{\pi^2 \varepsilon (1 - \bar{L} \varepsilon)}},$$

is crucial to prove the result.

**Remark 1.1.** *Let us give some remarks in order.*

1. *It is important to point out that the transmission conditions at the central node 0 are inspired by the recent papers [14, 16, 22, 28]. It is not the only possible choice, and the main motivation is that they guarantee uniqueness of the regular solutions of the (Bi - NLS) equation linearized around 0.*
2. *An important fact is that we are able to deal with the mix dispersion in the system (1.3), that is, with laplacian and bi-laplacian terms in the system. The laplacian term gives us an extra difficulty to deal with the adjoint system associated to (1.3). Precisely, if we remove the term  $\partial_x^2$  in (1.3) and deal only with the fourth order Schrödinger equation with the boundary conditions (1.4) we can use two different constants  $\alpha_j$  and  $\beta_j$  in the traces of the boundary conditions.*

<sup>1</sup>This method was introduced by V. Komornik in [26].

3. We are able to control  $N + 1$  edges with  $N$ -boundary controls, however, we do not have the sharp conditions on the lengths  $l_j$ . Moreover, the time of control  $T > T_{\min}$  is not sharp, but we get an explicit constant in the observability inequality. In this way, these two problems are open.
4. Finally, with respect to control results for the system (1.1) in a bounded domain, in a recent work, Ammari and Bouzidi [5], treated the exact controllability properties for the linear fourth-order Schrödinger equation with variable parameters and boundary conditions in an interval  $(0, l)$ , namely

$$(1.10) \quad \begin{cases} i\rho(x)\partial_t y = -\partial_x^2(\sigma(x)\partial_x^2 y) + \partial_x(q(x)\partial_x y)_x, & (t, x) \in (0, T) \times (0, \ell) \\ y(t, 0) = \partial_x y(t, 0) = y(t, \ell) = 0, \quad \partial_x y(t, \ell) = f(t), & t \in (0, T) \\ y(0, x) = y^0(x), x \in (0, \ell) \end{cases}$$

where  $f$  is a control that acts at the right  $x = \ell$ , the function  $y^0$  is the initial condition and the variable coefficients  $\rho$ ,  $\sigma$  and  $q$  satisfy some appropriate assumptions. Precisely, using the nonharmonic Fourier series, the authors showed spectral properties for the system (1.10). With these properties in hand, they are able to prove an observability inequality for the adjoint system, which is a consequence of Ingham's inequality due to Beurling. The results presented in [5] corroborate, in a bounded context, those presented in Theorem 1.1. The main difference of the result proved in [5] with the result in Theorem 1.1 can be seen in two ways: The results presented in [5] are in a subset of  $\mathbb{R}$  and one uses Ingham's inequality to prove the observability inequality and, consequently, the exact controllability property. In contrast, our result is in a compact star graphs structure and the observability inequality is shown using the multiplier method, which helps us to achieve exact controllability property.

To end our introduction, we present the outline of the manuscript. Section 2 is related with the well-posedness results for the system (1.3)-(1.4) and its adjoint. In Section 3, we give a rigorous proof of observability inequality, and with this in hand, we are able to prove Theorem 1.1. In Appendix A we present key lemmas using Morawetz multipliers which are crucial to prove the main result of the paper.

## 2. WELL-POSEDNESS RESULTS

**2.1. Linear system.** In this section we consider the following linear model

$$(2.1) \quad \begin{cases} i\partial_t u_j + \partial_x^2 u_j - \partial_x^4 u_j = 0, & (t, x) \in (0, T) \times I_j, \quad j = 1, 2, \dots, N \\ u_j(0, x) = u_{j0}(x), & x \in I_j, \quad j = 1, 2, \dots, N \end{cases}$$

with the boundary conditions

$$(2.2) \quad \begin{cases} u_1(-l_1, t) = \partial_x u_1(-l_1, t) = 0 \\ u_j(l_j, t) = \partial_x u_j(l_j, t) = 0, & j \in \{2, 3, \dots, N\}, \\ u_1(0, t) = \alpha_j u_j(0, t), & j \in \{2, 3, \dots, N\}, \\ \partial_x u_1(0, t) = \sum_{j=2}^N \frac{\partial_x u_j(0, t)}{\alpha_j} \\ \partial_x^2 u_1(0) = \alpha_j \partial_x^2 u_j(0, t), & j \in \{2, 3, \dots, N\}, \\ \partial_x^3 u_1(0, t) = \sum_{j=2}^N \frac{\partial_x^3 u_j(0, t)}{\alpha_j}. \end{cases}$$

Additionally, from now on we use the notation introduced in the introduction of the manuscript.

Let us consider the differential operator

$$A : u = (u_1, \dots, u_{N+1}) \in \mathcal{D}(A) \subset L^2(\mathcal{G}) \mapsto i\partial_x^2 u - i\partial_x^4 u \in L^2(\mathcal{R})$$

with domain defined by

$$(2.3) \quad D(A) := \left\{ u \in \prod_{j=1}^N H^4(I_j) \cap V : \partial_x u_1(0) = \sum_{j=2}^N \frac{\partial_x u_j(0)}{\alpha_j}, \quad \partial_x^3 u_1(0) = \sum_{j=2}^N \frac{\partial_x^3 u_j(0)}{\alpha_j} \right\},$$

where

$$(2.4) \quad V = \left\{ u \in \prod_{j=1}^N H^2(I_j) : u_1(-l_1) = \partial_x u_1(-l_1) = u_j(l_j) = \partial_x u_j(l_j) = 0, \right. \\ \left. u_1(0) = \alpha_j u_j(0), \partial_x^2 u_1(0) = \alpha_j \partial_x^2 u_j(0), \quad j \in \{2, 3, \dots, N\} \right\}$$

Then we can rewrite the homogeneous linear system (2.1)-(2.2) takes the form

$$(2.5) \quad \begin{cases} u_t(t) = Au(t), & t > 0 \\ u(0) = u_0 \in L^2(\mathcal{G}). \end{cases}$$

The following proposition guarantees some properties for the operator  $A$ . Precisely, the following holds.

**Proposition 2.1.** *The operator  $A : D(A) \subset L^2(\mathcal{G}) \rightarrow L^2(\mathcal{G})$  is self-adjoint in  $L^2(\mathcal{G})$ .*

*Proof.* Let us first to prove that  $A$  is a symmetric operator. To do this let  $u$  and  $v$  in  $D(A)$ . Then, by approximating  $u$  and  $v$  by  $C^4(\mathcal{G})$  functions, integrating by parts and using the boundary conditions (2.3) and (2.4) we have that

$$\begin{aligned} (Au, v)_{L^2(\mathcal{G})} &= \operatorname{Re} \int_{-l_1}^0 (Au)_1(x) \overline{v_1(x)} dx + \operatorname{Re} \sum_{j=2}^N \int_0^{l_j} (Au)_j(x) \overline{v_j(x)} dx \\ &= \operatorname{Re} \int_{-l_1}^0 (i\partial_x^2 u_1 - i\partial_x^4 u_1) \overline{v_1(x)} dx + \operatorname{Re} \sum_{j=2}^N \int_0^{l_j} (i\partial_x^2 u_j - i\partial_x^4 u_j)(x) \overline{v_j(x)} dx \\ &= \operatorname{Re} \int_{-l_1}^0 u_1(i\partial_x^2 \overline{v_1} - i\partial_x^4 \overline{v_1}) dx + \operatorname{Re} \sum_{j=2}^N \int_0^{l_j} u_j(i\partial_x^2 \overline{v_j} - i\partial_x^4 \overline{v_j}) dx \\ &\quad + \operatorname{Re} i \sum_{j=1}^N [\partial_x u_j \overline{v_j} - u_j \partial_x \overline{v_j} - \partial_x^3 u_j \overline{v_j} + \partial_x^2 u_j \partial_x \overline{v_j} - \partial_x u_j \partial_x^2 \overline{v_j} + u_j \partial_x^3 \overline{v_j}]_{\partial \mathcal{G}} \\ &= (u, Av)_{L^2(\mathcal{G})}, \quad \forall u, v \in D(A), \end{aligned}$$

that is,  $A$  is symmetric. It is not hard to see that  $D(A^*) = D(A)$ , so  $A$  is self-adjoint. This finishes the proof.  $\square$

By using semigroup theory,  $A$  generates a strongly continuous unitary group on  $L^2(\mathcal{G})$ , and for any  $u_0 = (u_{10}, u_{20}, \dots, u_{N0}) \in L^2(\mathcal{G})$  there exists a unique mild solution  $u \in C([0, T]; L^2(\mathcal{G}))$  of (2.5). Furthermore, if  $u_0 \in D(A)$ , then (2.5) has a classical solution satisfying  $u \in C([0, T]; D(A)) \cap C^1([0, T]; L^2(\mathcal{G}))$ . Summarizing, we have the following result.

**Proposition 2.2.** *Let  $u_0 = (u_{10}, u_{20}, \dots, u_{N0}) \in H_0^k(\mathcal{G})$ , for  $k \in \{0, 1, 2, 3, 4\}$ . Then the linear system (2.1) with boundary conditions (2.2) has a unique solution  $u$  on the space  $C([0, T]; H_0^k(\mathcal{G}))$ . In particular, for  $k = 4$  we get a classical solution and for the other cases ( $k \in \{0, 1, 2, 3\}$ ) the solution is a mild solution.*

**2.2. Adjoint system.** Now, we deal with the adjoint system associated to (2.1)-(2.2). As the operator  $A = i\partial_x^2 - i\partial_x^4$  is self adjoint (see Proposition 2.1) the adjoint system is defined as follows

$$(2.6) \quad \begin{cases} i\partial_t v_j + \partial_x^2 v_j - \partial_x^4 v_j = 0, & (t, x) \in (0, T) \times I_j, \quad j = 1, 2, \dots, N \\ v_j(T, x) = v_{jT}(x), & x \in I_j, \quad j = 1, 2, \dots, N \end{cases}$$

with the boundary conditions

$$(2.7) \quad \begin{cases} v_1(-l_1, t) = \partial_x v_1(-l_1, t) = 0 \\ v_j(l_j, t) = \partial_x v_j(l_j, t) = 0, & j \in \{2, 3, \dots, N\}, \\ v_1(0, t) = \alpha_j v_j(0, t), & j \in \{2, 3, \dots, N\}, \\ \partial_x v_1(0, t) = \sum_{j=2}^N \frac{\partial_x v_j(0, t)}{\alpha_j} \\ \partial_x^2 v_1(0) = \alpha_j \partial_x^2 v_j(0, t), & j \in \{2, 3, \dots, N\}, \\ \partial_x^3 v_1(0, t) = \sum_{j=2}^N \frac{\partial_x^3 v_j(0, t)}{\alpha_j}. \end{cases}$$

Also, as  $A = A^*$  we have that  $D(A^*) = D(A)$  and the proof of well-posedness is the same that in Proposition 2.2.

### 3. EXACT BOUNDARY CONTROLLABILITY

This section is devoted to the analysis of the exact controllability property for the linear system corresponding to (1.3) with boundary control (1.4). Here, we will present the answer for the control problem presented in the introduction of this work. First, let us present two definitions that will be important for the rest of the work.

**Definition 3.1.** Let  $T > 0$ , The system (1.3)-(1.4) is exactly controllable in time  $T$  if for any initial and final data  $u_0, u_T \in H^{-2}(\mathcal{G})$  there exist control functions  $h_j \in L^2(0, T)$ ,  $j \in \{2, 3, \dots, N\}$ , such that solution  $u$  of (1.3)-(1.4) on the tree shaped network of  $N + 1$  edges satisfies (1.5). In addition, when  $u_T = 0$  we said that the system (1.3)-(1.4) is null controllable in time  $T$ .

Now on, consider the transposition solution to (1.3)-(1.4), with  $h_1(t) = 0$ , which is given by the following.

**Definition 3.2.** We say  $u \in L^\infty(0, T; H^{-2}(\mathcal{G}))$  is solution of (1.3)-(1.4), with  $h_1(t) = 0$ , in the transposition sense if and only if

$$\sum_{j=1}^N \left( \int_0^T \langle u_j(t), f_j(t) \rangle dt + i \langle u_j(0), v_j(0) \rangle \right) + \sum_{j=2}^N \left( \int_{\mathcal{G}} h_j \partial_x^2 \bar{v}_j dx \right) = 0,$$

for every  $f \in L^2(0, T; H_0^2(\mathcal{G}))$ , where  $v(x, t)$  is the mild solution to the problem (2.6)-(2.7) on the space  $C([0, T]; H_0^2(\mathcal{G}))$ , with  $v(x, T) = 0$ , obtained in Proposition 2.2. Here,  $\langle \cdot, \cdot \rangle$  means the duality between the spaces  $H^{-2}(\mathcal{G})$  and  $H_0^2(\mathcal{G})$ .

With this in hand, the following lemma gives an equivalent condition for the exact controllability property.

**Lemma 3.1.** Let  $u_T \in H^{-2}(\mathcal{G})$ . Then, there exist controls  $h_j(t) \in L^2(0, T)$ , for  $j = 2, \dots, N$ , such that the solution  $u(x, t)$  of (1.3)-(1.4), with  $h_1(t) = 0$ , satisfies (1.5) if and only if

$$(3.1) \quad i \sum_{j=1}^N \int_{I_j} \langle u_j(T), \bar{v}_j(T) \rangle dx = \sum_{j=2}^N \int_0^T h_j(t) \partial_x^2 v_j(l_j, t) dt,$$

where  $v$  is solution of (2.6)-(2.7), with initial data  $v(x, T) = v(T)$ .

*Proof.* Relation (3.1) is obtained multiplying (1.3)-(1.4), with  $h_1(t) = 0$ , by the solution  $v$  of (2.6)-(2.7) and integrating by parts on  $\mathcal{G} \times (0, T)$ .  $\square$



**3.1. Observability inequality.** A fundamental role will be played by the following observability result, which together with Lemma 3.1 give us Theorem 1.1.

**Proposition 3.1.** *Let  $l_j > 0$  for any  $j \in \{1, \dots, N+1\}$  satisfying (1.6) and assume that (1.9) holds. There exists a positive constant  $T_{\min}$  such that if  $T > T_{\min}$ , then the following inequality holds*

$$(3.2) \quad \|v(x, T)\|_{H_0^2(\mathcal{G})}^2 \leq C \sum_{j=2}^N \|\partial_x^2 v_j(l_j, t)\|_{L^2(0, T)}^2$$

for any  $v = (v_1, v_2, \dots, v_{N+1})$  solution of (2.6)-(2.7) with final condition  $v_T = (v_1^T, v_2^T, \dots, v_{N+1}^T) \in H_0^2(\mathcal{G})$  and for a positive constant  $C > 0$ .

*Proof.* Firstly, taking  $f = 0$  and choosing  $q(x, t) = 1$  in (A.3), we get that

$$\begin{aligned} -\frac{Im}{2} \int_{\mathcal{G}} v \overline{\partial_x v} \Big|_0^T dx + \frac{Im}{2} \int_0^T v \overline{\partial_t v} \Big|_{\partial \mathcal{G}} dt + \frac{1}{2} \int_0^T |\partial_x v|^2 \Big|_{\partial \mathcal{G}} dt + \frac{1}{2} \int_0^T |\partial_x^2 v|^2 \Big|_{\partial \mathcal{G}} dt \\ - Re \int_0^T [\partial_x^3 v \overline{\partial_x v}]_{\partial \mathcal{G}} dt = 0, \end{aligned}$$

or equivalently,

$$\begin{aligned} 0 = & -\frac{Im}{2} \int_{\mathcal{G}} v \overline{\partial_x v} \Big|_0^T dx + \frac{Im}{2} \int_0^T \left( [v_1 \overline{\partial_t v_1}]_{-l_1}^0 + \left[ \sum_{j=2}^N v_j \overline{\partial_t v_j} \right]_0^{l_j} \right) dt \\ & + \frac{1}{2} \int_0^T \left( [|\partial_x v_1|^2]_{-l_1}^0 + \left[ \sum_{j=2}^N |\partial_x v_j|^2 \right]_0^{l_j} \right) dt + \frac{1}{2} \int_0^T \left( [|\partial_x^2 v_1|^2]_{-l_1}^0 + \left[ \sum_{j=2}^N |\partial_x^2 v_j|^2 \right]_0^{l_j} \right) dt \\ & - Re \int_0^T \left( [\partial_x^3 v_1 \overline{\partial_x v_1}]_{-l_1}^0 + \left[ \sum_{j=2}^N \partial_x^3 v_j \overline{\partial_x v_j} \right]_0^{l_j} \right) dt. \end{aligned}$$

By using the boundary conditions (2.7), it follows that

$$\begin{aligned} 0 = & -\frac{Im}{2} \int_{\mathcal{G}} v \overline{\partial_x v} \Big|_0^T dx + \frac{Im}{2} \int_0^T \left( v_1(0) \overline{\partial_t v_1}(0) - \sum_{j=2}^N v_j(0) \overline{\partial_t v_j}(0) \right) dt \\ & + \frac{1}{2} \int_0^T \left( |\partial_x v_1(0)|^2 - \sum_{j=2}^N |\partial_x v_j(0)|^2 \right) dt \\ & + \frac{1}{2} \int_0^T \left( |\partial_x^2 v_1(0)|^2 - |\partial_x^2 v_1(-l_1)|^2 + \sum_{j=2}^N (|\partial_x^2 v_j(l_j)|^2 - |\partial_x^2 v_j(0)|^2) \right) dt \\ & - Re \int_0^T \left( \partial_x^3 v_1(0) \overline{\partial_x v_1}(0) - \sum_{j=2}^N \partial_x^3 v_j(0) \overline{\partial_x v_j}(0) \right) dt. \end{aligned}$$

Once again, due to the boundary conditions (2.7) and relations (1.9), we have that

$$\begin{aligned} \int_0^T |\partial_x^2 v_1(-l_1)|^2 dt = & -Im \int_{\mathcal{G}} v \overline{\partial_x v} \Big|_0^T dx + \int_0^T \sum_{j=2}^N |\partial_x^2 v_j(l_j)|^2 dt \\ (3.3) \quad & + \int_0^T \left( |\partial_x v_1(0)|^2 - \sum_{j=2}^N |\partial_x v_j(0)|^2 \right) dt + \int_0^T \left( |\partial_x^2 v_1(0)|^2 - \sum_{j=2}^N |\partial_x^2 v_j(0)|^2 \right) dt. \end{aligned}$$



Thanks to relations (1.9), we deduce that

$$\begin{aligned} & \int_0^T \left( |\partial_x v_1(0)|^2 - \sum_{j=2}^N |\partial_x v_j(0)|^2 \right) dt = \frac{1}{2} \int_0^T \left( \left| \sum_{j=2}^N \frac{\partial_x v_j(0)}{\alpha_j} \right|^2 - \sum_{j=2}^N |\partial_x v_j(0)|^2 \right) dt \\ & \leq \frac{1}{2} \int_0^T \left( (N-1) \sum_{j=2}^N \left| \frac{\partial_x v_j(0)}{\alpha_j} \right|^2 - \sum_{j=2}^N |\partial_x v_j(0)|^2 \right) dt = \frac{1}{2} \int_0^T \sum_{j=2}^N |\partial_x v_j(0)|^2 \left( \frac{N-1}{\alpha_j^2} - 1 \right) dt \leq 0 \end{aligned}$$

and

$$\begin{aligned} & \frac{1}{2} \int_0^T \left( |\partial_x^2 v_1(0)|^2 - \sum_{j=2}^N |\partial_x^2 v_j(0)|^2 \right) dt = \frac{1}{2} \int_0^T \left( |\partial_x^2 v_1(0)|^2 - \sum_{j=2}^N \left| \frac{\partial_x^2 v_1(0)}{\alpha_j} \right|^2 \right) dt \\ & = \frac{1}{2} \int_0^T |\partial_x^2 v_1(0)|^2 \left( 1 - \sum_{j=2}^N \frac{1}{\alpha_j^2} \right) dt = 0. \end{aligned}$$

Thus, previous calculations ensure that

$$(3.4) \quad \int_0^T |\partial_x^2 v_1(-l_1)|^2 dt \leq \operatorname{Im} \int_{\mathcal{G}} v \overline{\partial_x v} \Big|_0^T dx + \int_0^T \sum_{j=2}^N |\partial_x^2 v_j(l_j)|^2 dt.$$

Now, choosing  $q(x, t) = x$  in (A.3), by using the boundary conditions (2.7) and taking  $f = 0$ , we get

$$\begin{aligned} 2 \int_Q |\partial_x v|^2 dx dt + 4 \int_Q |\partial_x^2 v|^2 dx dt &= \int_0^T \left[ \partial_x^2 v|^2 x \right]_{\partial \mathcal{G}} dt - \operatorname{Im} \int_{\mathcal{G}} v \overline{\partial_x^2 v} \Big|_0^T dx \\ &= l_1 \int_0^T |\partial_x^2 v_1(-l_1)|^2 dt + \int_0^T \sum_{j=2}^N |\partial_x^2 v_j(l_j)|^2 l_j dt - \operatorname{Im} \int_{\mathcal{G}} v \overline{\partial_x^2 v} \Big|_0^T dx. \end{aligned}$$

From inequality (3.4), it yields that

$$\begin{aligned} 2 \int_Q |\partial_x v|^2 dx dt + 4 \int_Q |\partial_x^2 v|^2 dx dt &\leq l_1 \operatorname{Im} \int_{\mathcal{G}} v \overline{\partial_x v} \Big|_0^T dx + l_1 \int_0^T \sum_{j=2}^N |\partial_x^2 v_j(l_j)|^2 dt \\ &\quad + \int_0^T \sum_{j=2}^N |\partial_x^2 v_j(l_j)|^2 l_j dt - \operatorname{Im} \int_{\mathcal{G}} v \overline{\partial_x^2 v} \Big|_0^T dx, \end{aligned}$$

hence

$$(3.5) \quad 2 \int_Q |\partial_x v|^2 dx dt + 4 \int_Q |\partial_x^2 v|^2 dx dt \leq \operatorname{Im} \int_{\mathcal{G}} v \overline{\partial_x v} (l_1 - x) \Big|_0^T dx + (\bar{L} + l_1) \int_0^T \sum_{j=2}^N |\partial_x^2 v_j(l_j)|^2 dt,$$

where  $\bar{L}$  is defined by (1.7).

Now, we are in position to prove (3.2). Thanks to (3.5), follows that

$$\begin{aligned} (3.6) \quad & 2 \int_Q (|\partial_x v|^2 + |\partial_x^2 v|^2) dx dt \leq (\bar{L} + l_1) \int_0^T \sum_{j=2}^N |\partial_x^2 v_j(l_j)|^2 dt + \left| \int_{\mathcal{G}} v \overline{\partial_x v} (l_1 - x) \Big|_0^T dx \right| \\ & \leq L \int_0^T \sum_{j=2}^N |\partial_x^2 v_j(l_j)|^2 dt + \int_{\mathcal{G}} |v(T)| |\overline{\partial_x v(T)}| |l_1 - x| dx \\ & \quad + \int_{\mathcal{G}} |v(0)| |\overline{\partial_x v(0)}| |l_1 - x| dx. \end{aligned}$$

As we have the conservation laws for solutions of (A.2), that is, (A.10) is satisfied, so by using it on the left hand side of (3.6), yields that

$$(3.7) \quad 2 \int_0^T (\|\partial_x v(T)\|_{L^2(\mathcal{G})}^2 + \|\partial_x^2 v(T)\|_{L^2(\mathcal{G})}^2) dt \leq L \int_0^T \sum_{j=2}^N |\partial_x^2 v_j(l_j)|^2 dt \\ + \bar{L} \left( \int_{\mathcal{G}} |v(T)| |\overline{\partial_x v(T)}| dx + \int_{\mathcal{G}} |v(0)| |\overline{\partial_x v(0)}| dx \right).$$

Applying Young inequality in (3.7), with  $\varepsilon > 0$  satisfying (1.8), we deduce that

$$(3.8) \quad 2T \left( \|\partial_x v(T)\|_{L^2(\mathcal{G})}^2 + \|\partial_x^2 v(T)\|_{L^2(\mathcal{G})}^2 \right) \leq L \int_0^T \sum_{j=2}^N |\partial_x^2 v_j(l_j)|^2 dt \\ + \bar{L} \left( \frac{1}{\varepsilon T} \int_{\mathcal{G}} |v(T)|^2 dx + \varepsilon T \int_{\mathcal{G}} |\overline{\partial_x v(T)}|^2 dx + \frac{1}{\varepsilon T} \int_{\mathcal{G}} |v(0)|^2 dx + \varepsilon T \int_{\mathcal{G}} |\overline{\partial_x v(0)}|^2 dx \right).$$

Therefore, we have due to (3.8) and, again using the conservation law, the following estimate

$$2T(1 - \bar{L}\varepsilon) \left( \|\partial_x v(T)\|_{L^2(\mathcal{G})}^2 + \|\partial_x^2 v(T)\|_{L^2(\mathcal{G})}^2 \right) \leq L \int_0^T \sum_{j=2}^N |\partial_x^2 v_j(l_j)|^2 dt + \frac{2\bar{L}}{\varepsilon T} \|v(T)\|_{L^2(\mathcal{G})}^2.$$

From relation (1.2), we have that

$$2(1 - \bar{L}\varepsilon)T \left( \|\partial_x v(T)\|_{L^2(\mathcal{G})}^2 + \|\partial_x^2 v(T)\|_{L^2(\mathcal{G})}^2 \right) \leq L \int_0^T \sum_{j=2}^N |\partial_x^2 v_j(l_j)|^2 dt \\ + \frac{2M}{\varepsilon T} \left( \frac{L^2}{\pi^2} + 1 \right) (\|\partial_x v(T)\|_{L^2(\mathcal{G})}^2 + \|\partial_x^2 v(T)\|_{L^2(\mathcal{G})}^2).$$

Equivalently, we get that

$$2\bar{L} \left[ \left( \frac{1}{\bar{L}} - \varepsilon \right) T - \frac{1}{\varepsilon T} \left( \frac{L^2}{\pi^2} + 1 \right) \right] (\|\partial_x v(T)\|_{L^2(\mathcal{G})}^2 + \|\partial_x^2 v(T)\|_{L^2(\mathcal{G})}^2) \leq L \int_0^T \sum_{j=2}^N |\partial_x^2 v_j(l_j)|^2 dt.$$

Note that the conditions (1.6), (1.7) and (1.8) imply that

$$K = \left[ \left( \frac{1}{\bar{L}} - \varepsilon \right) T - \frac{1}{\varepsilon T} \left( \frac{L^2}{\pi^2} + 1 \right) \right] > 0$$

Thus, again using (1.2), we achieved the observability inequality (3.2).  $\square$

**3.2. Proof of Theorem 1.1.** Notice that Theorem 1.1 is a consequence of the observability inequality (3.2). In fact, without loss of generality, pick  $u_0 = 0$  on  $\mathcal{G}$ . Define  $\Gamma$  the linear and bounded map  $\Gamma : H_0^2(\mathcal{G}) \longrightarrow H^{-2}(\mathcal{G})$  by

$$(3.9) \quad \Gamma(v(\cdot, T)) = \langle u(\cdot, T), \bar{v}(\cdot, T) \rangle,$$

where  $v = v(x, t)$  is solution of (2.6)-(2.7), with initial data  $v(x, T) = v(T)$ ,  $\langle \cdot, \cdot \rangle$  means the duality between the spaces  $H^{-2}(\mathcal{G})$  and  $H_0^2(\mathcal{G})$ ,  $u = u(x, t)$  is solution of (1.3)-(1.4), with  $h_1(t) = 0$  and

$$(3.10) \quad h_j(t) = \partial_x^2 v_j(l_j, t),$$

for  $j = 2, \dots, N$ .

According to Lemma 3.1 and Proposition 3.1, we obtain

$$\langle \Gamma(v(T)), v(T) \rangle = \sum_{j=2}^N \|h_j(t)\|_{L^2(0, T)}^2 \geq C^{-1} \|v(T)\|_{H_0^2(\mathcal{G})}^2.$$

Thus, by the Lax-Milgram theorem,  $\Gamma$  is invertible. Consequently, for given  $u(T) \in H^{-2}(\mathcal{G})$ , we can define  $v(T) := \Gamma^{-1}(u(T))$  which one solves (2.6)-(2.7). Then, if  $h_j(t)$ , for  $j = 2, \dots, N$  is defined

by (3.10), the corresponding solution  $u$  of the system (1.3)-(1.4), satisfies (1.5) and so, Theorem 1.1 holds.

## APPENDIX A. AUXILIARY LEMMAS

**A.1. Morawetz multipliers.** This section is dedicated to establishing fundamental identities by the multipliers method, which will be presented in two lemmas. For  $f \in L^2(0, T; H_0^2(\mathcal{G}))$ , let us consider the following system

$$(A.1) \quad \begin{cases} i\partial_t u_j + \partial_x^2 u_j - \partial_x^4 u_j = f, & (t, x) \in (0, T) \times I_j, \quad j = 1, 2, \dots, N \\ u_j(0, x) = u_{j0}(x), & x \in I_j, \quad j = 1, 2, \dots, N \end{cases}$$

with the boundary conditions

$$(A.2) \quad \begin{cases} u_1(-l_1, t) = \partial_x u_1(-l_1, t) = 0 \\ u_j(l_j, t) = \partial_x u_j(l_j, t) = 0, \quad j \in \{2, 3, \dots, N\}, \\ u_1(0, t) = \alpha_j u_j(0, t), \quad j \in \{2, 3, \dots, N\}, \\ \partial_x u_1(0, t) = \sum_{j=2}^N \frac{\partial_x u_j(0, t)}{\alpha_j} \\ \partial_x^2 u_1(0) = \alpha_j \partial_x^2 u_j(0, t), \quad j \in \{2, 3, \dots, N\}, \\ \partial_x^3 u_1(0, t) = \sum_{j=2}^N \frac{\partial_x^3 u_j(0, t)}{\alpha_j}. \end{cases}$$

The first lemma gives us an identity which will help us to prove the main result of this article.

**Lemma A.1.** *Let  $q = q(x, t) \in C^4(\bar{\mathcal{G}} \times (0, T), \mathbb{R})$  with  $\bar{\mathcal{G}}$  being the closed set of  $\mathcal{G}$ . For every solution of (A.2)-(A.1) with  $f \in \mathcal{D}(\mathcal{G})$  and  $u_0 \in \mathcal{D}(\mathcal{G})$ , the following identity holds:*

$$(A.3) \quad \begin{aligned} & \frac{Im}{2} \int_Q u \overline{\partial_x u} \partial_t q dx dt - \int_Q |\partial_x u|^2 \partial_x q dx dt - 2 \int_Q |\partial_x^2 u|^2 \partial_x q dx dt - \frac{Re}{2} \int_Q \partial_x u \bar{u} \partial_x^2 q dx dt \\ & + \frac{3}{2} \int_Q |\partial_x u|^2 \partial_x^3 q dx dt + \frac{Re}{2} \int_Q \partial_x u \bar{u} \partial_x^4 q dx dt - \frac{Im}{2} \int_Q u \overline{\partial_x u} q \Big|_0^T dx + \frac{1}{2} \int_0^T |\partial_x^2 u|^2 q \Big|_{\partial \mathcal{G}} dt \\ & + \frac{Im}{2} \int_0^T u \overline{\partial_t u} q \Big|_{\partial \mathcal{G}} dt + \frac{1}{2} \int_0^T |\partial_x u|^2 q \Big|_{\partial \mathcal{G}} dt + \frac{Re}{2} \int_0^T \partial_x u \bar{u} \partial_x q \Big|_{\partial \mathcal{G}} dt \\ & + \int_0^T \left[ -|\partial_x u|^2 \partial_x^2 q + \frac{3}{2} Re(\partial_x^2 u \overline{\partial_x u}) \partial_x q - Re(\partial_x^3 u \overline{\partial_x u}) q \right]_{\partial \mathcal{G}} dt \\ & + \int_0^T \left[ -\frac{Re}{2} (\partial_x u \bar{u}) \partial_x^3 q + \frac{Re}{2} (\partial_x^2 u \bar{u}) \partial_x^2 q - \frac{Re}{2} (\partial_x^3 u \bar{u}) \partial_x q \right] dt = \int_Q f(\overline{\partial_x u} q + \frac{1}{2} \bar{u} \partial_x q) dx dt, \end{aligned}$$

where  $Q := \mathcal{G} \times [0, T]$  and  $\partial \mathcal{G}$  is the boundary of  $\mathcal{G}$ .

*Proof.* Multiplying (A.1) by  $\overline{\partial_x u} q + \frac{1}{2} \bar{u} \partial_x q$ , we have that

$$\begin{aligned} & \int_Q i\partial_t u (\overline{\partial_x u} q + \frac{1}{2} \bar{u} \partial_x q) dx dt + \int_Q \partial_x^2 u (\overline{\partial_x u} q + \frac{1}{2} \bar{u} \partial_x q) dx dt - \int_Q \partial_x^4 u (\overline{\partial_x u} q + \frac{1}{2} \bar{u} \partial_x q) dx dt \\ & - \int_Q f (\overline{\partial_x u} q + \frac{1}{2} \bar{u} \partial_x q) dx dt := I_1 + I_2 - I_3 - \int_Q f (\overline{\partial_x u} q + \frac{1}{2} \bar{u} \partial_x q) dx dt. \end{aligned}$$

Now, we split the proof in three steps.

**Step 1.** Analysis of  $I_1$ .

Integrating by parts, several times, on  $Q$ , we get

$$\begin{aligned}
I_1 &= -i \int_Q u \overline{\partial_t \partial_x u} q dx dt - i \int_Q u \overline{\partial_x u} \partial_t q dx dt + i \left[ \int_{\mathcal{G}} u \overline{\partial_x u} q \right]_0^T dx + \frac{i}{2} \int_Q \partial_x u \overline{\partial_t u} q dx dt + \frac{i}{2} \int_Q u \overline{\partial_t \partial_x u} q dx dt \\
&\quad - \frac{i}{2} \left[ \int_0^T u \overline{\partial_t u} q \right]_{\partial \mathcal{G}} dt + \frac{i}{2} \int_Q \partial_x u \overline{u} \partial_t q dx dt + \frac{i}{2} \int_Q u \overline{\partial_x u} \partial_t q dx dt - \frac{i}{2} \left[ \int_0^T u \overline{u} \partial_t q \right]_{\partial \mathcal{G}} dt + \frac{i}{2} \left[ \int_{\mathcal{G}} u \overline{u} \partial_x q \right]_0^T dx \\
&= -\frac{i}{2} \int_Q u \overline{\partial_t \partial_x u} q dx dt - \frac{i}{2} \int_Q u \overline{\partial_x u} \partial_t q dx dt - \frac{i}{2} \int_Q \partial_t \partial_x u \overline{u} q dx dt + i \left[ \int_{\mathcal{G}} u \overline{\partial_x u} q \right]_0^T dx - \frac{i}{2} \left[ \int_0^T u \overline{\partial_t u} q \right]_{\partial \mathcal{G}} dt \\
&\quad - \frac{i}{2} \left[ \int_{\mathcal{G}} u \overline{\partial_x u} q \right]_0^T dx - \frac{i}{2} \left[ \int_{\mathcal{G}} u \overline{u} \partial_x q \right]_0^T dx + \frac{i}{2} [u \overline{u} q]_0^T|_{\partial \mathcal{G}} - \frac{i}{2} \left[ \int_0^T u \overline{u} \partial_t q \right]_{\partial \mathcal{G}} dt + \frac{i}{2} \left[ \int_{\mathcal{G}} u \overline{u} \partial_x q \right]_0^T dx.
\end{aligned}$$

Thus, putting together the similar terms in the last equality, yields that

$$\begin{aligned}
(A.4) \quad I_1 &= -\frac{i}{2} \int_Q (u \overline{\partial_t \partial_x u} + \partial_t \partial_x u \overline{u}) q dx dt - \frac{i}{2} \int_Q u \overline{\partial_x u} \partial_t q dx dt + \frac{i}{2} \left[ \int_{\mathcal{G}} u \overline{\partial_x u} q \right]_0^T dx \\
&\quad - \frac{i}{2} \left[ \int_0^T u \overline{\partial_t u} q \right]_{\partial \mathcal{G}} dt + \frac{i}{2} [u \overline{u} q]_0^T|_{\partial \mathcal{G}} - \frac{i}{2} \left[ \int_0^T u \overline{u} \partial_t q \right]_{\partial \mathcal{G}} dt.
\end{aligned}$$

Finally, taking the real part of (A.4), we have

$$(A.5) \quad Re(I_1) = \frac{Im}{2} \int_Q u \overline{\partial_x u} \partial_t q dx dt - \frac{Im}{2} \left[ \int_{\mathcal{G}} u \overline{\partial_x u} q \right]_0^T dx + Re \left( -\frac{i}{2} \left[ \int_0^T u \overline{\partial_t u} q \right]_{\partial \mathcal{G}} dt \right).$$

**Step 2.** Analysis of the Laplacian integral  $I_2$ .

Integrating by parts, several times, on  $Q$  and taking the real part of  $I_2$ , follows that

$$\begin{aligned}
Re(I_2) &= \int_Q Re(\partial_x^2 u \overline{\partial_x u} q) dx dt + \frac{Re}{2} \left( \int_Q \partial_x^2 u \overline{u} \partial_x q dx dt \right) \\
&= -\frac{1}{2} \int_Q |\partial_x u|^2 \partial_x q dx dt + \frac{1}{2} \left[ \int_0^T |\partial_x u|^2 q \right]_{\partial \mathcal{G}} dt - \frac{1}{2} \int_Q |\partial_x u|^2 \partial_x q dx dt \\
&\quad - \frac{Re}{2} \left( \int_Q \partial_x u \overline{u} \partial_x^2 q dx dt \right) + \frac{Re}{2} \left( \left[ \int_0^T \partial_x u \overline{u} \partial_x q \right]_{\partial \mathcal{G}} dt \right).
\end{aligned}$$

Consequently,

$$\begin{aligned}
(A.6) \quad Re(I_2) &= - \int_Q |\partial_x u|^2 \partial_x q dx dt - \frac{Re}{2} \left( \int_Q \partial_x u \overline{u} \partial_x^2 q dx dt \right) \\
&\quad + \frac{1}{2} \left[ \int_0^T |\partial_x u|^2 q \right]_{\partial \mathcal{G}} dt + \frac{Re}{2} \left( \left[ \int_0^T \partial_x u \overline{u} \partial_x q \right]_{\partial \mathcal{G}} dt \right).
\end{aligned}$$

**Step 3.** Analysis of the bi-Laplacian integral  $I_3$ .

Integrating by parts on  $Q$  give us

$$\begin{aligned}
(A.7) \quad -I_3 &= \int_Q \partial_x^3 u \overline{\partial_x^2 u} q dx dt + \int_Q \partial_x^3 u \overline{\partial_x u} \partial_x q dx dt - \left[ \int_0^T \partial_x^3 u \overline{\partial_x u} q \right]_{\partial \mathcal{G}} dt \\
&\quad + \frac{1}{2} \int_Q \partial_x^3 u \overline{\partial_x u} \partial_x q dx dt + \frac{1}{2} \int_Q \partial_x^3 u \overline{u} \partial_x^2 q dx dt - \frac{1}{2} \left[ \int_0^T \partial_x^3 u \overline{u} \partial_x q \right]_{\partial \mathcal{G}} dt.
\end{aligned}$$

Taking the real part of (A.7) and integrating, again, by parts on  $Q$ , we have

$$\begin{aligned}
-Re(I_3) = & -\frac{1}{2} \int_Q |\partial_x^2 u|^2 \partial_x q dx dt + \frac{1}{2} \int_0^T \left[ \partial_x^2 u |^2 q \right]_{\partial \mathcal{G}} dt - \frac{3}{2} \int_Q |\partial_x^2 u|^2 \partial_x q dx dt + \int_Q |\partial_x u|^2 \partial_x^3 q dx dt \\
& - \int_0^T \left[ |\partial_x u|^2 \partial_x^2 q \right]_{\partial \mathcal{G}} dt + \frac{3Re}{2} \left( \int_0^T \left[ \partial_x^2 u \overline{\partial_x u} \partial_x q \right]_{\partial \mathcal{G}} dt \right) - Re \left( \int_0^T \left[ \partial_x^3 u \overline{\partial_x u} q \right]_{\partial \mathcal{G}} dt \right) \\
& + \frac{Re}{2} \left( \int_Q |\partial_x u|^2 \partial_x^3 q dx dt \right) + \frac{Re}{2} \left( \int_Q \partial_x u \overline{u} \partial_x^4 q dx dt \right) - \frac{Re}{2} \left( \int_0^T \left[ \partial_x u \overline{u} \partial_x^3 q \right]_{\partial \mathcal{G}} dt \right) \\
& + \frac{Re}{2} \left( \int_0^T \left[ \partial_x^2 u \overline{u} \partial_x^2 q \right]_{\partial \mathcal{G}} dt \right) - \frac{Re}{2} \left( \int_0^T \left[ \partial_x^3 u \overline{u} \partial_x q \right]_{\partial \mathcal{G}} dt \right).
\end{aligned}$$

The last inequality ensure that

$$\begin{aligned}
(A.8) \quad -Re(I_3) = & -2 \int_Q |\partial_x^2 u|^2 \partial_x q dx dt + \frac{3}{2} \int_Q |\partial_x u|^2 \partial_x^3 q dx dt + \frac{Re}{2} \left( \int_Q \partial_x u \overline{u} \partial_x^4 q dx dt \right) \\
& + \frac{1}{2} \int_0^T \left[ \partial_x^2 u |^2 q \right]_{\partial \mathcal{G}} dt - \int_0^T \left[ |\partial_x u|^2 \partial_x^2 q \right]_{\partial \mathcal{G}} dt + \frac{3Re}{2} \left( \int_0^T \left[ \partial_x^2 u \overline{\partial_x u} \partial_x q \right]_{\partial \mathcal{G}} dt \right) \\
& - Re \left( \int_0^T \left[ \partial_x^3 u \overline{\partial_x u} q \right]_{\partial \mathcal{G}} dt \right) - \frac{Re}{2} \left( \int_0^T \left[ \partial_x u \overline{u} \partial_x^3 q \right]_{\partial \mathcal{G}} dt \right) \\
& + \frac{Re}{2} \left( \int_0^T \left[ \partial_x^2 u \overline{u} \partial_x^2 q \right]_{\partial \mathcal{G}} dt \right) - \frac{Re}{2} \left( \int_0^T \left[ \partial_x^3 u \overline{u} \partial_x q \right]_{\partial \mathcal{G}} dt \right).
\end{aligned}$$

Finally, taking into account (A.5), (A.6) and (A.8), we have

$$Re(I_1) + Re(I_2) - Re(I_3) = \int_Q f(\overline{\partial_x u} q + \frac{1}{2} \overline{u} \partial_x q) dx dt,$$

then (A.3) holds.  $\square$

**A.2. Conservation laws.** The final result of the appendix gives us the conservation laws for the solutions of (A.1)-(A.2), with  $f = 0$ . More precisely, the result is the following one.

**Lemma A.2.** *For any positive time  $t$ , the solution  $u$  of (A.1)-(A.2), with  $f = 0$ , satisfies*

$$(A.9) \quad \|u(t)\|_{L^2(\mathcal{G})} = \|u(0)\|_{L^2(\mathcal{G})}$$

and

$$(A.10) \quad \|\partial_x u(t)\|_{L^2(\mathcal{G})} + \|\partial_x^2 u(t)\|_{L^2(\mathcal{G})} = \|\partial_x u(0)\|_{L^2(\mathcal{G})} + \|\partial_x^2 u(0)\|_{L^2(\mathcal{G})}.$$

Additionally, we have

$$(A.11) \quad \|u(t)\|_{L^2(\mathcal{G})} + \|\partial_x u(t)\|_{L^2(\mathcal{G})} + \|\partial_x^2 u(t)\|_{L^2(\mathcal{G})} = \|u(0)\|_{L^2(\mathcal{G})} + \|\partial_x u(0)\|_{L^2(\mathcal{G})} + \|\partial_x^2 u(0)\|_{L^2(\mathcal{G})}.$$

*Proof.* Multiplying system (A.1)-(A.2), with  $f = 0$ , by  $i\overline{u}$  and integrating in  $\mathcal{G}$ , we get

$$(A.12) \quad 0 = - \int_{\mathcal{G}} \partial_t u_j \overline{u} dx + i \int_{\mathcal{G}} \partial_x^2 u_j \overline{u} dx - i \int_{\mathcal{G}} \partial_x^4 u_j \overline{u} dx = L_1 + L_2 + L_3.$$

We are now looking for the integral  $L_2 + L_3$ . Let us, first, rewrite these quantities as follows

$$(A.13) \quad L_2 + L_3 = i \sum_{j=1}^N \left( \int_{I_j} \partial_x^2 u_j \overline{u}_j dx - \int_{I_j} \partial_x^4 u_j \overline{u}_j dx \right).$$

Integrating (A.13) by parts and taking the real part of  $L_2 + L_3$ , we get that

$$\begin{aligned}
 (A.14) \quad \operatorname{Re}(L_2 + L_3) &= \operatorname{Re} \left( i \sum_{j=1}^N \left( - \int_{I_j} |\partial_x u_j|^2 dx + \partial_x u_j \bar{u}_j \right) \right) \\
 &\quad + \operatorname{Re} \left( i \sum_{j=1}^N \left( - \int_{I_j} |\partial_x^2 u_j|^2 dx + \partial_x^2 u_j \partial_x \bar{u}_j \right) - \partial_x^3 u_j \bar{u}_j \right) \\
 &= \operatorname{Re} i \left( \partial_x u_1 \bar{u}_1 + \partial_x^2 u_1 \partial_x \bar{u}_1 - \partial_x^3 u_1 \bar{u}_1 \right) \Big|_{-l_1}^0 \\
 &\quad + \operatorname{Re} i \sum_{j=2}^N \left( \partial_x u_j \bar{u}_j + \partial_x^2 u_j \partial_x \bar{u}_j - \partial_x^3 u_j \bar{u}_j \right) \Big|_0^{l_j}.
 \end{aligned}$$

By using the boundary conditions (A.2), we have

$$\begin{aligned}
 (A.15) \quad \operatorname{Re}(L_2 + L_3) &= \operatorname{Re} i \left( \partial_x u_1(0) \bar{u}_1(0) + \partial_x^2 u_1(0) \partial_x \bar{u}_1(0) - \partial_x^3 u_1(0) \bar{u}_1(0) \right) \\
 &\quad + \operatorname{Re} i \sum_{j=2}^N \left( -\partial_x u_j(0) \bar{u}_j(0) - \partial_x^2 u_j(0) \partial_x \bar{u}_j(0) + \partial_x^3 u_j(0) \bar{u}_j(0) \right) = 0.
 \end{aligned}$$

Thus, replacing (A.15) in (A.12), (A.9) holds.

We will prove (A.10). Multiplying (A.1), with  $f = 0$ , by  $\overline{\partial_t u}$ , integrating on  $\mathcal{G}$  and taking the real part give us

$$(A.16) \quad \operatorname{Re} \left( i \int_{\mathcal{G}} |\partial_t u|^2 dx \right) + \operatorname{Re} \left( \int_{\mathcal{G}} \partial_x^2 u \overline{\partial_t u} dx \right) - \operatorname{Re} \left( \int_{\mathcal{G}} \partial_x^4 u \overline{\partial_t u} dx \right) = 0.$$

Integrating (A.16) by parts on  $\mathcal{G}$  and using the boundary conditions (A.2), yields that

$$\begin{aligned}
 (A.17) \quad \frac{1}{2} \frac{\partial}{\partial t} \int_{\mathcal{G}} (|\partial_x u|^2 + |\partial_x^2 u|^2) dx &= -\operatorname{Re} \left[ \partial_x^2 u_1(0) \partial_x \partial_t \bar{u}_1(0) + \partial_x u_1(0) \partial_t \bar{u}_1(0) - \partial_x^3 u_1(0) \partial_t \bar{u}_1(0) \right] \\
 &\quad + \sum_{j=2}^N \left( -\partial_x^2 u_j(0) \partial_x \partial_t \bar{u}_j(0) - \partial_x u_j(0) \partial_t \bar{u}_j(0) + \partial_x^3 u_j(0) \partial_t \bar{u}_j(0) \right).
 \end{aligned}$$

The boundary condition give us that the right hand side of (A.17) is zero, that is

$$\frac{1}{2} \frac{\partial}{\partial t} \int_{\mathcal{G}} (|\partial_x u|^2 + |\partial_x^2 u|^2) dx = 0,$$

which implies (A.10). Finally, adding (A.9) and (A.10), we have (A.11).  $\square$

**Acknowledgments.** The authors wish to thank the referee for his/her valuable comments which improved this paper. Capistrano-Filho was supported by CNPq grants 408181/2018-4 and 307808/2021-1, CAPES grants 88881.311964/2018-01, 88881.520205/2020-01, MATHAMSUD 21-MATH-03 and Propesqi (UFPE). Cavalcante was supported by CNPq 310271/2021-5 and CAPES-MATHAMSUD 88887.368708/2019-00. Gallego was supported by MATHAMSUD 21-MATH-03 and the 100.000 Strong in the Americas Innovation Fund. This work was carried out during visits of the authors to the Universidade Federal de Alagoas, Universidade Federal de Pernambuco and Universidad Nacional de Manizales. The authors would like to thank the universities for its hospitality.

## REFERENCES

- [1] R. Adami, C. Cacciapuoti, D. Finco and D. Noja, *Variational properties and orbital stability of standing waves for NLS equation on a star graph*, J. Differential Equations, 257, 3738–3777 (2014).
- [2] R. Adami, C. Cacciapuoti, D. Finco and D. Noja, *Stable standing waves for a NLS on star graphs as local minimizers of the constrained energy*, J. Differential Equations, 260, 7397–7415 (2016).

- [3] J. Angulo and N. Goloshchapova, *On the orbital instability of excited states for the NLS equation with the  $\delta$ -interaction on a star graph*, Discrete Contin. Dyn. Syst. A., 38, no. 10, 5039–5066 (2018).
- [4] J. Angulo and N. Goloshchapova, *Extension theory approach in the stability of the standing waves for the NLS equation with point interactions on a star graph*, Adv. Differential Equations 23, no. 11-12, 793–846 (2018).
- [5] K. Ammari and H. Bouzidi, *Exact boundary controllability of the linear Biharmonic Schrödinger equation with variable coefficients*, arXiv:2112.15196 [math.AP] (2021).
- [6] K. Ammari and E. Crépeau, *Feedback Stabilization and Boundary Controllability of the Korteweg-de Vries Equation on a Star-Shaped Network*, SIAM J. Control Optim., 56, (3), 1620–1639 (2018).
- [7] K. Ammari and E. Crépeau, *Well-posedness and stabilization of the Benjamin-Bona-Mahony equation on star-shaped networks*, Systems & Control Letters, 127, 39–43 (2019).
- [8] L. Baudouin and M. Yamamoto, *Inverse problem on a tree-shaped network: unified approach for uniqueness*, Applicable Analysis, 94, (11), 2370–2395 (2015).
- [9] M. Ben-Artzi, H. Koch and J.-C Saut, *Dispersion estimates for fourth order Schrödinger equations*, C. R. Acad. Sci. Paris Sér. I Math, 330, 87–92 (2000).
- [10] G. Berkolaiko and P. Kuchment, *Introduction to Quantum Graphs*, Mathematical Surveys and Monographs, 186, Amer. Math. Soc., Providence, RI, (2013).
- [11] J. Blank, P. Exner and M. Havlicek, *Hilbert Space Operators in Quantum Physics*, 2nd edition, Theoretical and Mathematical Physics, Springer, New York, 2008.
- [12] J. L. Bona and R. C. Cascaval, *Nonlinear dispersive waves on trees*, Can. Appl. Math. Q. 16, no. 1, 1–18 (2008).
- [13] R. Burioni, D. Cassi, M. Rasetti, P. Sodano and A. Vezzani, *Bose-Einstein condensation on inhomogeneous complex networks*, J. Phys. B: At. Mol. Opt. Phys., 34, 4697–4710 (2001).
- [14] R. de A. Capistrano-Filho, M. Cavalcante and F. A. Gallego, *Lower regularity solutions of the biharmonic Schrödinger equation in a quarter plane*, Pacific Journal of Mathematics, 309-1, 35-70 (2020).
- [15] R. A. Capistrano-Filho, M. Cavalcante and F. A. Gallego, *Forcing operators on star graphs applied for the cubic fourth order Schrödinger equation*, Discrete & Continuous Dynamical Systems - B, doi: 10.3934/dcdsb.2021190.
- [16] M. Cavalcante, *The Korteweg-de Vries equation on a metric star graph*, Z. Angew. Math. Phys. 69: 124 (2018).
- [17] E. Cerpa, E. Crépeau and C. Moreno, *On the boundary controllability of the Korteweg-de Vries equation on a star-shaped network*, IMA Journal of Mathematical Control and Information, Vol. 37, No. 1, 226–240 (2020).
- [18] E. Cerpa, E. Crépeau, J. Valein, *Boundary controllability of the Korteweg-de Vries equation on a tree-shaped network*, Evol. Equ. Control Theory, Vol. 9, No. 3, 673–692 (2020).
- [19] A. Duca, *Global Exact Controllability of Bilinear Quantum Systems on Compact Graphs and Energetic Controllability*, SIAM J. Control Optim., 58(6), 3092–3129 (2020).
- [20] A. Duca, *Bilinear quantum systems on compact graphs: Well-posedness and global exact controllability*, Automatica, Vol. 123, 109324 (2021).
- [21] G. Fibich, B. Ilan and G. Papanicolaou, *Self-focusing with fourth-order dispersion*, SIAM J. Appl. Math. 62, 1437–1462 (2002).
- [22] F. Gregorio and D. Mugnolo, *Bi-Laplacians on graphs and networks*, J. Evol. Equ. 20, 191–232 (2020).
- [23] L. I. Ignat, A. F. Pazoto and L. Rosier, *Inverse problem for the heat equation and the Schrödinger equation on a tree*, Inverse Problems, 28 (1), (2011).
- [24] V. I. Karpman, *Stabilization of soliton instabilities by higher-order dispersion: fourth order nonlinear Schrödinger-type equations*, Phys. Rev. E, 53, 1336–1339 (1996).
- [25] V. I. Karpman and A. G. Shagalov, *Stability of soliton described by nonlinear Schrödinger type equations with higher-order dispersion*, Physica D, 144, 194–210 (2000).
- [26] V. Komornik, *Exact Controllability and Stabilization. The Multiplier Method Collection*, RMA vol 36 (Paris Masson) (1994).
- [27] D. Mugnolo, *Mathematical Technology of Networks*, Bielefeld, Springer Proceedings in Mathematics & Statistics 128, 2015.
- [28] D. Mugnolo, D. Noja and C. Seifert, *Airy-type evolution equations on star graphs*, Analysis & PDE. 11 (7), 1625–1652 (2018).
- [29] D. Mugnolo and J. F. Rault, *Construction of exact travelling waves for the Benjamin-Bona-Mahony equation on networks*, Bull. Belg. Math. Soc. Simon Stevin. 21, 415–436 (2014).
- [30] T. Tsutsumi, *Strichartz estimates for Schrödinger equation of fourth order with periodic boundary condition*, Kyoto university, 11pp, (2014).

*Departamento de Matemática, Universidade Federal de Pernambuco (UFPE), 50740-545, Recife (PE), Brazil.*  
*Email address: roberto.capistranofilho@ufpe.br*

*Instituto de Matemática, Universidade Federal de Alagoas (UFAL), Maceió (AL), Brazil*  
*Email address: marcio.melo@im.ufal.br*

*Departamento de Matemáticas y Estadística, Universidad Nacional de Colombia (UNAL), Cra 27 No. 64-60, 170003, Manizales, Colombia*  
*Email address: fagallegor@unal.edu.co*