

CONTROLS INSENSITIZING THE NORM OF SOLUTION OF A SCHRÖDINGER TYPE SYSTEM WITH MIXED DISPERSION

ROBERTO DE A. CAPISTRANO-FILHO* AND THIAGO YUKIO TANAKA

ABSTRACT. The main goal of this manuscript is to prove the existence of insensitizing controls for the fourth-order dispersive nonlinear Schrödinger equation with cubic nonlinearity. To obtain the main result we prove a null controllability property for a coupled fourth-order Schrödinger system of cascade type with zero order coupling which is equivalent to the insensitizing control problem. Precisely, by means of new Carleman estimates, we first obtain a null controllability result for the linearized system around zero, then the null controllability for the nonlinear case is extended using an inverse mapping theorem.

1. INTRODUCTION

1.1. Setting of the problem. Inspired by the difficulties in finding data in applications of distributed systems, Lions [27] introduced the topic of insensitizing controls. Precisely, this kind of problem deals with the existence of controls making a functional (depending on the solution) insensible to small perturbations of the initial data. Considering some particular functional, it has been proven that this problem is equivalent to control properties of cascade systems [4, 19].

The insensitivity can be defined in two different ways: An approximate problem or an exact problem. Approximate insensitivity is equivalent to approximate controllability of the cascade system, while exact insensitivity is equivalent to its null controllability. Before giving the reader more details about it and a state of the art related to these problems, let us introduce the model we want to study.

In this article, we address the exact insensitizing problem for the cubic fourth-order Schrödinger equation with mixed dispersion, the so-called fourth-order nonlinear Schrödinger system (4NLS)

$$(1.1) \quad iu_t + u_{xx} - u_{xxxx} = \lambda|u|^2u,$$

where $x, t \in \mathbb{R}$ and $u(x, t)$ is a complex-valued function. Equation (1.1) has been introduced by Karpman [23] and Karpman and Shagalov [24] to take into account the role of small fourth-order dispersion terms in the propagation of intense laser beams in a bulk medium with Kerr nonlinearity. Equation (1.1) arises in many scientific fields such as quantum mechanics, nonlinear optics and plasma physics, and has been intensively studied with fruitful references (see [3, 23, 30] and references therein).

To introduce our problem, consider $\Omega := (0, L) \subset \mathbb{R}$ be an interval and assume that $T > 0$. We will use the following notations $Q_T = \Omega \times (0, T)$, $\Sigma = \partial\Omega \times (0, T)$ and $\omega_T = \omega \times (0, T)$, where $\omega \subset \Omega$ is the so-called *control domain*. Let us consider the system

$$(1.2) \quad \begin{cases} iu_t + u_{xx} - u_{xxxx} - \zeta|u|^2u = f + 1_\omega h, & \text{in } Q_T, \\ u(t, 0) = u(t, L) = u_x(t, 0) = u_x(t, L) = 0, & \text{on } t \in (0, T), \\ u(0, x) = u_0(x) + \tau \hat{u}_0(x), & \text{in } \Omega, \end{cases}$$

where $\zeta \in \mathbb{R}$, τ is an unknown (small) real number, h stands for the control, u is the state function and $\hat{u}_0(x)$ is an unknown function.

2020 *Mathematics Subject Classification.* 35Q55, 93C20, 93B05, 93B07, 93C41.

Key words and phrases. Fourth-order Schrödinger system, Carleman estimates, Null controllability, Insensitizing controls.

*Corresponding author: roberto.capistranofilho@ufpe.br.

Let $J : \mathbb{R} \times L^2(q_T) \rightarrow \mathbb{R}$ be a functional (called *sentinel functional*) defined by

$$(1.3) \quad J(\tau, h) := \frac{1}{2} \iint_{\mathcal{O}_T} |u(x, t; \tau, h)|^2 dx dt,$$

where $u = u(x, t; \tau, h)$ is the corresponding solution of (1.2) associated to τ , h is the control function and $\mathcal{O}_T = \mathcal{O} \times (0, T)$, where \mathcal{O} is the so-called *observation domain*. Thus, our objective can be expressed in the definition below.

Definition 1 (Insensitizing controls). *Let $u_0 \in L^2(\Omega)$ and $f \in L^2(Q_T)$. We say that a control h insensitizes the functional J , associated with the solution $u(x, t; \tau, h)$ of (1.2), if*

$$(1.4) \quad \left. \frac{\partial}{\partial \tau} J(\tau, h) \right|_{\tau=0} = 0, \quad \forall \hat{u}_0 \in L^2(\Omega) \text{ with } \|\hat{u}_0\|_{L^2(\Omega)} = 1.$$

The definition (1.3) above can be seen as this: the sentinel does not detect (small) variations of the initial data u_0 provoked by the unknown (small) perturbation $\tau \hat{u}_0$ in the observation domain \mathcal{O} when the system evolve from a time $t = 0$ to a time $t = T$.

1.2. Insensitizing control problems for PDEs. As we have mentioned, the first time that insensitizing problem was approached was in the early of 90's by Lions [27, 28], where the author studied second and fourth order parabolic equations in limited domains considering a functional with the local L^2 -norm of the solution of a system with null initial data and where the control domain (located internally) intersect the observation domain (the set where we want to analyze the functional).

Since then, variations of this problem have been considered, i.e., to find controls which turn a functional depending on the solution (or some derivative) insensitive to small perturbations depending on the initial data. We will give a brief state of the art to the reader, precisely, we will present a sample of the insensitizing problems for partial differential equations (PDEs) and some control results to the system (1.1).

The first mathematical results concerned the insensitivity of the L^2 -norm of the solution restricted to a subdomain, called the observatory. In [11], the author proves that insensitizing control problems cannot be solved for every initial data. Additionally, de Tereza [14] used a global Carleman estimate approach to get the existence of exact insensitizing controls for a semilinear heat equation. Still with respect to semilinear heat equation, Bodart *et al.* [4, 5] proved the existence of insensitizing controls for this system with nonlinear boundary Fourier conditions.

Concerning to the variations of the sentinel functional, in [19], the author consider a functional involving the gradient of the state for a linear heat system, in the same way, Guerrero [21] treated the case of the sentinel with the curl of the solution of a Stokes system. With regard to wave equation, Alabau-Boussouira [1], showed the exact controllability, by a reduced number of controls, of coupled cascade systems of PDE's and the existence of exact insensitizing controls for the scalar wave equation. She gave a necessary and sufficient condition for the observability of abstract-coupled cascade hyperbolic systems by a single observation, the observation operator being either bounded or unbounded.

A variation of the (exact) control strategy was presented in [4], where the authors considered an approximated insensitizing problem (called ϵ -insensitizing control) for a nonhomogeneous heat equation. We observe that by smoothing the control strategy it was possible to prove in [13] positive results where the control domain and observation region do not intersect each other. Also in [10] the author proved insensitizing control results on unbounded domains, in [29] the authors treated insensitizing controls for both linear and semilinear heat equation but with a partially unknown domain, finally see [32] for the semilinear parabolic equation with dynamic boundary conditions. It is important to point out that in [22] the second author also treated it with a gradient type sentinel associated with the solutions of a nonlinear Ginzburg-Landau equation.

With respect to the structure/type of the equations/systems many variations were considered. In [8] the author treated insensitizing controls for the Boussinesq systems and in [7] the authors proved a result for a phase field system. In [17] it is considered a Cahn-Hilliard equation of

fourth order with superlinear nonlinearity and in [16] the authors proved insensitizing (exact and approximated) controls for a large-scale ocean circulation model.

To finalize this small sample of the state of the art, we cite Bodart *et al.* [5, 6] that studied systems with nonlinearities with certain superlinear growth and nonlinear terms depending on the state and its gradient. For a dispersive problem, we can cite Kumar and Chong [31] that worked with the KdV-Burgers equation. Finally, let us mention a recent work due Lopez-Garcia *et al.* [12]. In this work, the authors presented a control problem for a cascade system of two linear N -dimensional Schrödinger equations. They address the problem of null controllability by means of a control supported in a region not satisfying the classical geometrical control condition. The proof is based on the application of a Carleman estimate with degenerate weights to each one of the equations and a careful analysis of the system in order to prove null controllability with only one control force.

We caution that the literature is vast and one can see the references cited previously for the existence of the insensitizing controls for other types of PDEs.

1.3. Main results. The main goal of this paper is to close the gap that was missing when discussing the insensitizing control for the Schrödinger type equation. Here, we present the insensitizing control for the Schrödinger type equation with mixed dispersion. Precisely, we are interested in proving the existence of a control h which insensitizes the functional J defined by (1.3). The first result of this article can be read as follows.

Theorem 1.1. *Assume that $\omega \cap \mathcal{O} \neq \emptyset$ and $u_0 \equiv 0$. There exists a constant $C > 0$ and $\delta > 0$ such that for any f satisfying*

$$\|e^{C/t}f\|_{L^2(Q_T)} \leq \delta,$$

one can find a control $h(x, t) =: h \in L^2(Q_T)$ which insensitizes the functional J defined by (1.3), in the sense of Definition 1.

As mentioned at the beginning of this work, the existence of insensitizing controls for (1.2) can be defined equivalently by means of a null controllability problem for a cascade type system similar to the initial (1.2). In fact, the process consists in analysing the adjoint state of (1.2), more precisely by means of Definition 1 and since J is given by (1.3), the left-hand side of the second equation of (1.5) is the adjoint state of the derivative of (1.2) with respect to τ (with $\tau = 0$) and the right-hand side couples this equation with the localized state $1_{\mathcal{O}}u$. Then it is not difficult to see that condition (1.4) is equivalent to $v|_{t=0} \equiv 0$ in Ω^1 . So, in this spirit, due to the choice of J , we will reformulate our goal as a partial null controllability problem to the nonlinear system of cascade type associated to (1.2). In other words:

Problem A: *Can we find a control $h(x, t) = h \in L^2(Q_T)$ such that the solutions (u, v) of the following optimality coupled system*

$$(1.5) \quad \begin{cases} iu_t + u_{xx} - u_{xxxx} - \zeta|u|^2u = f + 1_{\omega}h, & \text{in } Q_T, \\ iv_t + v_{xx} - v_{xxxx} - \bar{\zeta}\bar{u}^2\bar{v} - 2\bar{\zeta}|u|^2v = 1_{\mathcal{O}}u, & \text{in } Q_T, \\ u(t, 0) = u(t, L) = v(t, 0) = v(t, L) = 0, & \text{on } t \in (0, T), \\ u_x(t, 0) = u_x(t, L) = v_x(t, 0) = v_x(t, L) = 0, & \text{on } t \in (0, T), \\ u(0, x) = u_0(x), \quad v(T, x) = 0, & \text{in } \Omega, \end{cases}$$

satisfies, in the time $t = 0$, $v|_{t=0} = 0$?

The answer to such a question motivates the next theorem, which is the main result of this paper.

Theorem 1.2. *Assume that $\omega \cap \mathcal{O} \neq \emptyset$ and the initial data $u_0 \equiv 0$. Then, there exists positive constants C and δ , depending on ω , Ω , \mathcal{O} , ζ and T , such that for any f satisfying*

$$\|e^{C/t}f\|_{L^2(Q_T)} \leq \delta,$$

¹See [4, 27] for a rigorous deduction of this fact and [33] for an explicit computation to obtain the cascade system with a general nonlinearity for a Ginzburg-Landau equation.

there exists a control $h \in L^2(q_T)$ such that the corresponding solution (u, v, h) of (1.5) satisfies $v|_{t=0} = 0$ in Ω .

Thus, to find controls insensitizing the functional J , that is, to prove Theorem 1.1, it is sufficient to prove the partial null controllability result given in Theorem 1.2. Therefore, from now on, we will concentrate on proving Theorem 1.2.

1.4. Heuristic and structure of the manuscript. Let us now explain the ideas to prove the results introduced in the last subsection. The main strategy adopted is based on duality arguments (see, e.g. [15, 26]). Roughly speaking, we prove suitable observability inequalities for the solutions of an adjoint system, where the main tool is a new Carleman estimate. This Carleman estimate with right-hand side in weight Sobolev spaces will be the key point to deal with the coupling terms of the linear system associated to (1.5).

In details, to prove Theorem 1.2 we will first prove a null controllability result for the linearized system associated to (1.5) around zero, which is given by

$$(1.6) \quad \begin{cases} iu_t + u_{xx} - u_{xxxx} = f^0 + 1_\omega h, & \text{in } Q_T, \\ iv_t + v_{xx} - v_{xxxx} = f^1 + 1_\mathcal{O} u, & \text{in } Q_T, \\ u(t, 0) = u(t, L) = v(t, 0) = v(t, L) = 0, & \text{on } t \in (0, T), \\ u_x(t, 0) = u_x(t, L) = v_x(t, 0) = v_x(t, L) = 0, & \text{on } t \in (0, T), \\ u(0, x) = u_0(x), v(T, x) = 0, & \text{in } \Omega. \end{cases}$$

Here, f^0 and f^1 are (small) source terms in appropriated L^p -weighted spaces. In order to prove it, we consider the adjoint system of (1.6), namely,

$$(1.7) \quad \begin{cases} i\varphi_t + \varphi_{xx} - \varphi_{xxxx} = 1_\mathcal{O} \psi + g^0, & \text{in } Q_T, \\ i\psi_t + \psi_{xx} - \psi_{xxxx} = g^1, & \text{in } Q_T, \\ \varphi(t, 0) = \varphi(t, L) = \varphi_x(t, 0) = \varphi_x(t, L) = 0, & \text{on } t \in (0, T), \\ \psi(t, 0) = \psi(t, L) = \psi_x(t, 0) = \psi_x(t, L) = 0, & \text{on } t \in (0, T), \\ \varphi(T, x) = 0, \psi(0, x) = \psi_0(x), & \text{in } \Omega. \end{cases}$$

With this in hand, we are able to prove an *observability inequality*, with aspect like the one below,

$$(1.8) \quad \iint_{Q_T} \rho_1(|\varphi|^2 + |\psi|^2) dx dt \leq C \iint_{\omega_T} \rho_2|\varphi|^2 dx dt + \iint_{Q_T} \rho_3(|g^0|^2 + |g^1|^2) dx dt,$$

where ρ_i , $i = 1, 2, 3$, are appropriate weights functions. Then, by duality approach, the desired partial null controllability property is a direct consequence of the (1.8) and can be read as follows.

Theorem 1.3. *Assuming that $\omega \cap \mathcal{O} \neq \emptyset$ and the initial data $u_0 \equiv 0$, there exists a positive constant C , depending on $\delta, \omega, \Omega, \mathcal{O}$ and T , such that for f^0 and f^1 , in a suitable weighted spaces, one can find a control h such that the associated solution (u, v) of (1.6) satisfies $v|_{t=0} \equiv 0$ in Ω .*

The last step is to use an inverse mapping theorem to extend the previous result to the nonlinear system.

Remarks 1. *Finally, the following comments are now given in order:*

1. *In the Definition 1, it is important to point out that the data u_0 will be taken in a convenient way, precisely, will be taken such that the functional J , given by (1.3), is well-defined.*
2. *On the Theorem 1.2, the smallness of f is related to the fact that we will apply a local inversion argument, that is, we first study this problem when a linearized form of equation (1.5) is considered and then we apply a local inversion mapping theorem.*
3. *We claim that Theorem 1.1 is equivalent to Theorem 1.2. In fact, considering (u, v) solution of (1.5) and using the boundary conditions of (1.5), a direct calculation leads us to the following*

$$(1.9) \quad \left. \frac{\partial}{\partial \tau} J(\tau, h) \right|_{\tau=0} = -\operatorname{Re} \int_{\Omega} i \hat{u}_0 v(0) dx.$$

Therefore, we can conclude that the left-hand side of (1.9) is zero, for all $\hat{u}_0 \in L^2(\Omega)$ with $\|\hat{u}_0\|_{L^2(\Omega)} = 1$, if and only if, $v(0) = 0$ in Ω .

4. It is worth mentioning that, in our work, we need a Carleman estimate with internal observation, differently from what was proven by Zheng [34]. Another interesting point is that on the Zheng's work, he proved the regularity of solution of the 4NLS in the class

$$C^1([0, T]; L^2(\Omega)) \cap C^0([0, T]; H^3(\Omega) \cap H_0^2(\Omega)),$$

which is also different in our case, we need more regular solutions (see Section A) to help us to use the inverse mapping theorem.

5. Finally, observe that our sentinel functional J is defined in the sense of L^2 -norm. If we want to insensitize a functional with a norm greater than L^2 , for example, $\partial_x^n u$, for $n \geq 1$, then we need a system coupled in the second equation of (1.5) in the form $\partial_x^n(1_{\mathcal{O}} \partial_x^n u)$, this means, the coupling has twice as many derivatives. More details about this kind of problem will be given in Section 4.

Our work is outlined in the following way: Section 2 is devoted to presenting a new Carleman estimate which will be the key to prove the main result of this manuscript. In the Section 3, we show the null controllability results, that is, the linear case (Theorem 1.3) and the nonlinear one (Theorem 1.2). Section 4, we present further comments and some open problems which seem to be of interest from the mathematical point of view. Finally, for completeness, at the end of this paper, we have an Appendix A about the existence of solutions for the systems considered here.

2. CARLEMAN ESTIMATES

In this section we prove a new Carleman estimate to the fourth order Schrödinger operator $\partial_x^4 u - \partial_x^2 u$. For the sake of simplicity, we will consider the operator $\mathcal{L} = \partial_x^4 u$, that is, only with the higher term. So, to derive this new Carleman, first, let us introduce the basic weight function $\eta(x) = (x - x_0)$ with $x_0 < 0$. Now, for $\lambda > 1$ and $\mu > 1$ we define the following

$$(2.1) \quad \theta = e^l, \quad \xi(t, x) = \frac{e^{3\mu\eta(x)}}{t(T-t)} \quad \text{and} \quad l(t, x) = \lambda \frac{e^{3\mu\eta(x)} - e^{5\mu|\eta|_\infty}}{t(T-t)}.$$

Our result will be derived from a previous result due to Zheng [34], which can be seen as follows.

Proposition 2.1. *There exists three constants $\mu_0 > 1$, $C_0 > 0$ and $C > 0$ such that for all $\mu > \mu_0$ and for all $\lambda \geq C_0(T + T^2)$,*

$$(2.2) \quad \begin{aligned} & \iint_{Q_T} (\lambda^7 \mu^8 \xi^7 \theta^2 |u|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |u_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |u_{xx}|^2 + \lambda \mu^2 \xi \theta^2 |u_{xxx}|^2) dx dt \\ & \leq C \left(\iint_{Q_T} |\theta Pu|^2 dx dt + \lambda^3 \mu^3 \int_0^T (\xi^3 \theta^2 |u_{xx}|^2)(t, L) dt + \lambda \mu \int_0^T (\xi \theta^2 |u_{xxx}|^2)(t, L) dt \right), \end{aligned}$$

where $Pu = i\partial_t u + \partial_x^4 u$.

With the previous theorem in hand, we are in position to prove a new Carleman estimate associated to the operator $\mathcal{L}u$. The result is stated in the following way.

Theorem 2.2. *Let $\omega, \mathcal{O} \subset \Omega$ be open subsets such that $\omega \cap \mathcal{O} \neq \emptyset$. Then, there exists a positive constant μ_1 , such that for any $\mu > \mu_1$, one can find two positive constants λ_1 and C depending on $\lambda, \mu, \Omega, \omega$ such that for any $\lambda > \lambda_1(T + T^2)$ the following estimate for φ and ψ of (2.4) holds*

$$(2.3) \quad \begin{aligned} & \iint_{Q_T} (\lambda^7 \mu^8 \xi^7 \theta^2 |\varphi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\varphi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\varphi_{xx}|^2 + \lambda \mu^2 \xi \theta^2 |\varphi_{xxx}|^2) dx dt \\ & + \iint_{Q_T} (\lambda^7 \mu^8 \xi^7 \theta^2 |\psi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\psi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\psi_{xx}|^2 + \lambda \mu^2 \xi \theta^2 |\psi_{xxx}|^2) dx dt \\ & \leq C \left(\iint_{Q_T} \theta^2 (|g^0|^2 + |g^1|^2) dx dt + \lambda \mu \iint_{\omega_T} \xi \theta^2 |\varphi|^2 dx dt \right). \end{aligned}$$

Before proving to prove this result, let us give the idea to derive (2.3). To do it, we split the proof in several steps. The first one, consists in applying the Carleman estimate given by Proposition 2.1 for (φ, ψ) solutions of

$$(2.4) \quad \begin{cases} i\varphi_t + \varphi_{xx} - \varphi_{xxxx} = 1_{\mathcal{O}}\psi + g^0, & \text{in } Q_T, \\ i\psi_t + \psi_{xx} - \psi_{xxxx} = g^1, & \text{in } Q_T, \\ \varphi(t, 0) = \varphi(t, L) = \varphi_x(t, 0) = \varphi_x(t, L) = 0, & \text{on } t \in (0, T), \\ \psi(t, 0) = \psi(t, L) = \psi_x(t, 0) = \psi_x(t, L) = 0, & \text{on } t \in (0, T), \\ \varphi(T, x) = 0, \psi(0, x) = \psi_0, & \text{in } \Omega. \end{cases}$$

The second step concern the estimate for a local integral term of ψ in terms of a local integral of φ and global integral terms of g^0, g^1, φ, ψ and smaller order terms of ψ . Finally, we will estimate integral terms on the border in terms of global integral of φ, ψ and smaller order integral terms.

Proof of Theorem 2.2. In what follows, remember that $\Omega \subset \mathbb{R}$ is a bounded domain whose boundary $\partial\Omega$ is regular enough. Consider $T > 0$, ω and \mathcal{O} to be two nonempty subsets of Ω . Additionally, as defined at the beginning of this work $Q_T = \Omega \times (0, T)$, $\omega_T = \omega \times (0, T)$, $\Sigma_T = \partial\Omega \times (0, T)$, $\mathcal{O}_T = \mathcal{O} \times (0, T)$ and denote by C a generic constant which can be different from one computation to another. Thus, let us split the proof in three steps.

Step 1: Applying Carleman estimates (2.2).

Thanks to (2.2) we have, for φ and ψ , solution of (2.4), that

$$(2.5) \quad \begin{aligned} & \iint_{Q_T} (\lambda^7 \mu^8 \xi^7 \theta^2 |\varphi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\varphi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\varphi_{xx}|^2 + \lambda \mu^2 \xi \theta^2 |\varphi_{xxx}|^2) dx dt \\ & \leq C \left(\iint_{Q_T} \theta^2 |\varphi_{xx}|^2 dx dt + \iint_{Q_T} \theta^2 |g^0|^2 dx dt + \iint_{\mathcal{O}_T} \theta^2 |\psi|^2 dx dt \right. \\ & \quad \left. + \lambda^3 \mu^3 \int_0^T (\xi^3 \theta^2 |\varphi_{xx}|^2)(t, L) dt + \lambda \mu \int_0^T (\xi \theta^2 |\varphi_{xxx}|^2)(t, L) dt \right). \end{aligned}$$

For λ, μ large enough, we obtain

$$(2.6) \quad \begin{aligned} & \iint_{Q_T} (\lambda^7 \mu^8 \xi^7 \theta^2 |\varphi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\varphi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\varphi_{xx}|^2 + \lambda \mu^2 \xi \theta^2 |\varphi_{xxx}|^2) dx dt \\ & \leq C \left(\iint_{Q_T} \theta^2 |g^0|^2 dx dt + \iint_{\mathcal{O}_T} \theta^2 |\psi|^2 dx dt + \lambda^3 \mu^3 \int_0^T (\xi^3 \theta^2 |\varphi_{xx}|^2)(t, L) dt \right. \\ & \quad \left. + \lambda \mu \int_0^T (\xi \theta^2 |\varphi_{xxx}|^2)(t, L) dt \right). \end{aligned}$$

Now, applying (2.2) for ψ , we get that

$$(2.7) \quad \begin{aligned} & \iint_{Q_T} (\lambda^7 \mu^8 \xi^7 \theta^2 |\psi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\psi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\psi_{xx}|^2 + \lambda \mu^2 \xi \theta^2 |\psi_{xxx}|^2) dx dt \\ & \leq C \left(\iint_{Q_T} \theta^2 |\psi_{xx}|^2 dx dt + \iint_{Q_T} \theta^2 |g^1|^2 dx dt + \iint_{\mathcal{O}_T} |\psi|^2 dx dt \right. \\ & \quad \left. + \lambda^3 \mu^3 \int_0^T (\xi^3 \theta^2 |\psi_{xx}|^2)(t, L) dt + \lambda \mu \int_0^T (\xi \theta^2 |\psi_{xxx}|^2)(t, L) dt \right). \end{aligned}$$

Again, by using (2.7) for λ, μ large enough we have the following:

$$(2.8) \quad \begin{aligned} & \iint_{Q_T} (\lambda^7 \mu^8 \xi^7 \theta^2 |\psi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\psi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\psi_{xx}|^2 + \lambda \mu^2 \xi \theta^2 |\psi_{xxx}|^2) dx dt \\ & \leq C \left(\iint_{Q_T} \theta^2 |g^1|^2 dx dt + \lambda^3 \mu^3 \int_0^T (\xi^3 \theta^2 |\psi_{xx}|^2)(t, L) dt + \lambda \mu \int_0^T (\xi \theta^2 |\psi_{xxx}|^2)(t, L) dt \right). \end{aligned}$$

Note that, putting together (2.6) and (2.8), we obtain the following estimate

$$\begin{aligned}
 & \iint_{Q_T} (\lambda^7 \mu^8 \xi^7 \theta^2 |\varphi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\varphi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\varphi_{xx}|^2 + \lambda \mu^2 \xi \theta^2 |\varphi_{xxx}|^2) dx dt \\
 & + \iint_{Q_T} (\lambda^7 \mu^8 \xi^7 \theta^2 |\psi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\psi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\psi_{xx}|^2 + \lambda \mu^2 \xi \theta^2 |\psi_{xxx}|^2) dx dt \\
 (2.9) \quad & \leq C \left(\iint_{Q_T} \theta^2 |g^0|^2 dx dt + \iint_{Q_T} \theta^2 |g^1|^2 dx dt + \iint_{\mathcal{O}_T} \theta^2 |\psi|^2 dx dt \right. \\
 & \left. + \lambda^3 \mu^3 \int_0^T (\xi^3 \theta^2 (|\varphi_{xx}|^2 + |\psi_x|^2))(t, L) dt + \lambda \mu \int_0^T (\xi \theta^2 (|\varphi_{xxx}|^2 + |\psi_{xxx}|^2))(t, L) dt \right).
 \end{aligned}$$

Step 2: Estimates for the local integral of ψ .

In this step, let us estimate the last term in the right-hand side of (2.9), that is, the local integral term of ψ in \mathcal{O}_T . Now, since $\omega \cap \mathcal{O} \neq \emptyset$, there exists $\tilde{\omega}_T \subset \omega \cap \mathcal{O}$. From now on, take a cut-off function $\eta \in C_0^\infty(\omega)$ such that $\eta \equiv 1$ in $\tilde{\omega}_T$. Observe that

$$\psi = -i\varphi_t + \varphi_{xx} - \varphi_{xxxx} - g^0, \quad \text{in } \mathcal{O}_T,$$

so

$$\begin{aligned}
 \iint_{\mathcal{O}_T} \theta^2 |\psi|^2 dx dt & \leq \iint_{\tilde{\omega}_T} \eta \theta^2 |\psi|^2 dx dt = \iint_{\tilde{\omega}_T} \eta \theta^2 \bar{\psi} \psi dx dt \\
 (2.10) \quad & = \operatorname{Re} \iint_{\tilde{\omega}_T} \eta \theta^2 \bar{\psi} (-i\varphi_t + \varphi_{xx} - \varphi_{xxxx} - g^0) dx dt \\
 & := \sum_{i=1}^4 \Psi_i,
 \end{aligned}$$

where Ψ_i , for $i = 1, 2, 3, 4$, are the integrals of the right-hand side of (2.10). We now estimate these terms. For Ψ_1 , integrating by parts in t we have that

$$\begin{aligned}
 \Psi_1 & = \operatorname{Re} \iint_{\tilde{\omega}_T} \eta \theta^2 \bar{\psi} (-i\varphi_t) dx dt = \operatorname{Re} \left((\eta \theta^2 \bar{\psi} (-i\varphi)) \Big|_0^T - \iint_{Q_T} (\eta \theta^2 \bar{\psi})_t (-i\varphi) dx dt \right) \\
 (2.11) \quad & = \operatorname{Re} \left(- \iint_{\tilde{\omega}_T} (\eta \theta^2)_t \bar{\psi} (-i\varphi) dx dt - \iint_{\tilde{\omega}_T} \eta \theta^2 \bar{\psi}_t (-i\varphi) dx dt \right).
 \end{aligned}$$

Since,

$$|(\theta^m)_t| = |(e^{ml})_t| = |e^{ml}(ml)_t| = |me^{ml}(l)_t| \leq mC\lambda T \xi^2 \theta^m,$$

then, for $m = 2$, we get that $|(\eta \theta^2)_t| \leq C\lambda \xi^2 \theta^2$, and by using Young inequality we obtain

$$\begin{aligned}
 (2.12) \quad \operatorname{Re} \left(- \iint_{\tilde{\omega}_T} (\eta \theta^2)_t \bar{\psi} (-i\varphi) dx dt \right) & \leq C \operatorname{Re} \left(\lambda \iint_{\tilde{\omega}_T} \xi^2 \theta^2 \bar{\psi} \varphi dx dt \right) \\
 & = \operatorname{Re} \left(\iint_{\tilde{\omega}_T} \left(\lambda^{\frac{7}{2}} \mu^4 \xi^{\frac{7}{2}} \theta \bar{\psi} \right) \left(C \lambda^{-\frac{5}{2}} \mu^{-4} \xi^{-\frac{3}{2}} \varphi \right) dx dt \right) \\
 & \leq \delta \lambda^7 \mu^8 \iint_{Q_T} \xi^7 \theta^2 |\psi|^2 dx dt + C \lambda^{-5} \mu^{-8} \iint_{\tilde{\omega}_T} \xi^{-3} |\varphi|^2 dx dt.
 \end{aligned}$$

Combining (2.11) with (2.12), we get

$$\begin{aligned}
 (2.13) \quad \Psi_1 & = \operatorname{Re} \left(- \iint_{\tilde{\omega}_T} \eta \theta^2 \bar{\psi}_t (-i\varphi) dx dt \right) \\
 & \quad + \delta \lambda^7 \mu^8 \iint_{Q_T} \xi^7 \theta^2 |\psi|^2 dx dt + C \lambda^{-5} \mu^{-8} \iint_{\tilde{\omega}_T} \xi^{-3} |\varphi|^2 dx dt.
 \end{aligned}$$

So, for δ small enough we can absorb the global integral term of (2.13) with the left-hand side of (2.9). Due to the boundary conditions for ψ and φ and since η has compact support on $\tilde{\omega}$ by integrating by parts for space variable, we obtain

$$\begin{aligned}
 \Psi_2 &= Re \iint_{\tilde{\omega}_T} \eta \theta^2 \bar{\psi} \varphi_{xx} dx dt = Re \left(\iint_{\tilde{\omega}_T} (\eta \theta^2 \bar{\psi})_{xx} \varphi dx dt \right) \\
 (2.14) \quad &= Re \left(\iint_{\tilde{\omega}_T} (\eta \theta^2)_{xx} \bar{\psi} \varphi dx dt + 2 \iint_{\tilde{\omega}_T} (\eta \theta^2)_x \bar{\psi}_x \varphi dx dt + \iint_{\tilde{\omega}_T} \eta \theta^2 \bar{\psi}_{xx} \varphi dx dt \right) \\
 &= \sum_{i=1}^3 \Psi_{2,i}.
 \end{aligned}$$

To bound each term of the right-hand side of (2.14) first observe that the following estimates, for the derivative in space of the weight function θ , holds true

$$|(\theta^2)_x| = \left| (e^{2l})_x \right| = \left| e^{2l} (2l)_x \right| = 2 |\theta^2 l_x| \leq C \lambda \mu \xi \theta^2$$

and

$$\begin{aligned}
 |(\theta^2)_{xx}| &= \left| (e^{2l})_{xx} \right| = \left| (e^{2l} (2l)_x)_x \right| = 2 |(\theta^2 l_x)_x| \\
 &\leq 4 |\theta^2 l_x^2| + 2 |\theta^2 l_{xx}| \leq 4 \theta^2 |l_x|^2 + 2 \theta^2 |l_{xx}| \\
 &\leq C \lambda^2 \mu^2 \xi^2 \theta^2 + C \lambda \mu \xi^2 \theta^2 \leq C \lambda^2 \mu^2 \xi^2 \theta^2.
 \end{aligned}$$

Therefore, it yields that

$$\begin{aligned}
 \Psi_{2,1} &= Re \left(\iint_{\tilde{\omega}_T} (\eta \theta^2)_{xx} \bar{\psi} \varphi dx dt \right) \leq Re \left(C \lambda^2 \mu^2 \iint_{\tilde{\omega}_T} \xi^2 \theta^2 \bar{\psi} \varphi dx dt \right) \\
 (2.15) \quad &= Re \left(\iint_{\tilde{\omega}_T} \left(\lambda^{\frac{7}{2}} \mu^4 \xi^{\frac{7}{2}} \theta \bar{\psi} \right) \left(C \lambda^{-\frac{3}{2}} \mu^{-2} \xi^{-\frac{3}{2}} \theta \varphi \right) dx dt \right) \\
 &\leq \delta \lambda^7 \mu^8 \iint_{Q_T} \xi^7 \theta^2 |\psi|^2 dx dt + C \lambda^{-3} \mu^{-4} \iint_{\tilde{\omega}_T} \xi^{-3} \theta^2 |\varphi|^2 dx dt
 \end{aligned}$$

and

$$\begin{aligned}
 \Psi_{2,2} &= 2 Re \left(\iint_{\tilde{\omega}_T} (\eta \theta^2)_x \bar{\psi}_x \varphi dx dt \right) \leq Re \left(C \lambda \mu \iint_{\tilde{\omega}_T} \xi \theta^2 \bar{\psi} \varphi dx dt \right) \\
 (2.16) \quad &= 2 Re \left(\iint_{\tilde{\omega}_T} \left(\lambda^{\frac{5}{2}} \mu^3 \xi^{\frac{5}{2}} \theta \bar{\psi}_x \right) \left(C \lambda^{-\frac{3}{2}} \mu^{-2} \xi^{-\frac{3}{2}} \theta \varphi \right) dx dt \right) \\
 &\leq \delta \lambda^5 \mu^6 \iint_{Q_T} \xi^5 \theta^2 |\psi_x|^2 dx dt + C \lambda^{-3} \mu^{-4} \iint_{\tilde{\omega}_T} \xi^{-3} \theta^2 |\varphi|^2 dx dt.
 \end{aligned}$$

Finally, $\Psi_{2,3}$ does not need to be estimated since we use it to obtain the equation for ψ . Now, combining (2.14), (2.15) and (2.16) we get

$$\begin{aligned}
 \Psi_2 &\leq C \lambda^{-3} \mu^{-4} \iint_{\tilde{\omega}_T} \xi^{-3} \theta^2 |\varphi|^2 dx dt \\
 (2.17) \quad &+ \delta \left(\lambda^7 \mu^8 \iint_{Q_T} \xi^7 \theta^2 |\psi|^2 dx dt + \lambda^7 \mu^8 \iint_{Q_T} \xi^7 \theta^2 |\psi|^2 dx dt + \lambda^7 \mu^8 \iint_{Q_T} \xi^7 \theta^2 |\psi_{xx}|^2 dx dt \right).
 \end{aligned}$$

For Ψ_3 we use the boundary conditions and integrate with respect to space variable four times to obtain

$$\begin{aligned}
 \Psi_3 &= \operatorname{Re} \iint_{\tilde{\omega}_T} \eta \theta^2 \bar{\psi} (-\varphi_{xxxx}) dx dt = \operatorname{Re} \left(\iint_{\tilde{\omega}_T} (\eta \theta^2 \bar{\psi})_{xxxx} (-\varphi) dx dt \right) \\
 &= \operatorname{Re} \left(\iint_{\tilde{\omega}_T} (\eta \theta^2)_{xxxx} \bar{\psi} \varphi dx dt + 4 \iint_{\tilde{\omega}_T} (\eta \theta^2)_{xxx} \bar{\psi}_x \varphi dx dt + 6 \iint_{\tilde{\omega}_T} (\eta \theta^2)_{xx} \bar{\psi}_{xx} \varphi dx dt \right) \\
 &\quad + \operatorname{Re} \left(4 \iint_{\tilde{\omega}_T} (\eta \theta^2)_x \bar{\psi}_{xxx} \varphi dx dt + \iint_{\tilde{\omega}_T} \eta \theta^2 (-\bar{\psi}_{xxxx}) \varphi dx dt \right) \\
 &= \sum_{i=1}^5 \Psi_{3,i}.
 \end{aligned} \tag{2.18}$$

Now our task is to estimate these terms. Observe that we have the following estimates in space variable for k -th order derivative in space variable for the weight function θ ,

$$|(\theta^2)_{kx}| \leq 2^k C \left(\lambda^k \mu^k \xi^k \right) \theta^2,$$

thus

$$\begin{aligned}
 \Psi_{3,1} &= \operatorname{Re} \left(\iint_{\tilde{\omega}_T} (\eta \theta^2)_{xxxx} \bar{\psi} \varphi dx dt \right) \\
 &\leq \operatorname{Re} \left(C \lambda^4 \mu^4 \iint_{\tilde{\omega}_T} \xi^4 \theta^2 \bar{\psi} \varphi dx dt \right) \\
 &= \operatorname{Re} \left(\iint_{\tilde{\omega}_T} \left(\lambda^{\frac{7}{2}} \mu^4 \xi^{\frac{7}{2}} \theta \bar{\psi} \right) \left(C \lambda^{\frac{1}{2}} \xi^{\frac{1}{2}} \theta \varphi \right) dx dt \right) \\
 &\leq \delta \lambda^7 \mu^8 \iint_{Q_T} \xi^7 \theta^2 |\psi|^2 dx dt + C \lambda \iint_{\tilde{\omega}_T} \xi \theta^2 |\varphi|^2 dx dt,
 \end{aligned} \tag{2.19}$$

$$\begin{aligned}
 \Psi_{3,2} &= \operatorname{Re} \left(4 \iint_{\tilde{\omega}_T} (\eta \theta^2)_{xxx} \bar{\psi}_x \varphi dx dt \right) \\
 &\leq \operatorname{Re} \lambda^3 \mu^3 \left(\iint_{\tilde{\omega}_T} \xi^3 \theta^2 \bar{\psi}_x \varphi dx dt \right) \\
 &= \operatorname{Re} \left(\iint_{\tilde{\omega}_T} \left(\lambda^{\frac{5}{2}} \mu^3 \xi^{\frac{5}{2}} \theta \bar{\psi}_x \right) \left(C \lambda^{\frac{1}{2}} \xi^{\frac{1}{2}} \theta \varphi \right) dx dt \right) \\
 &\leq \delta \lambda^5 \mu^6 \iint_{Q_T} \xi^5 \theta^2 |\psi_x|^2 dx dt + C \lambda \iint_{\tilde{\omega}_T} \xi \theta^2 |\varphi|^2 dx dt,
 \end{aligned} \tag{2.20}$$

$$\begin{aligned}
 \Psi_{3,3} &= \operatorname{Re} \left(6 \iint_{\tilde{\omega}_T} (\eta \theta^2)_{xx} \bar{\psi}_{xx} \varphi dx dt \right) \\
 &\leq \operatorname{Re} \left(C \lambda^2 \mu^2 \iint_{\tilde{\omega}_T} \xi^2 \theta^2 \bar{\psi}_{xx} \varphi dx dt \right) \\
 &= \operatorname{Re} \left(\iint_{\tilde{\omega}_T} \left(\lambda^{\frac{3}{2}} \mu^2 \xi^{\frac{3}{2}} \theta \bar{\psi}_{xx} \right) \left(C \lambda^{\frac{1}{2}} \xi^{\frac{1}{2}} \theta \varphi \right) dx dt \right) \\
 &\leq \delta \lambda^3 \mu^4 \iint_{Q_T} \xi^3 \theta^2 |\psi_{xx}|^2 dx dt + C \lambda \iint_{\tilde{\omega}_T} \xi \theta^2 |\varphi|^2 dx dt
 \end{aligned} \tag{2.21}$$

and

$$\begin{aligned}
\Psi_{3,4} &= \operatorname{Re} \left(4 \iint_{\tilde{\omega}_T} (\eta \theta^2)_x \bar{\psi}_{xxx} \varphi dx dt \right) \\
&\leq \operatorname{Re} \left(C \lambda \mu \iint_{\tilde{\omega}_T} \xi \theta^2 \bar{\psi}_{xxx} \varphi dx dt \right) \\
(2.22) \quad &= \operatorname{Re} \left(\iint_{\tilde{\omega}_T} \left(\lambda^{\frac{1}{2}} \mu \xi^{\frac{1}{2}} \theta \bar{\psi}_{xxx} \right) \left(C \lambda^{\frac{1}{2}} \xi^{\frac{1}{2}} \theta \varphi \right) dx dt \right) \\
&\leq \delta \lambda \mu^2 \iint_{Q_T} \xi \theta^2 |\psi_{xxx}|^2 dx dt + C \lambda \iint_{\tilde{\omega}_T} \xi \theta^2 |\varphi|^2 dx dt.
\end{aligned}$$

We do not estimate $\Psi_{3,5}$ since we use this term to obtain the equation for ψ . By putting (2.19), (2.20), (2.21) and (2.22) in (2.18), we conclude

$$\begin{aligned}
\Psi_3 &\leq \operatorname{Re} \left(\iint_{\tilde{\omega}_T} \eta \theta^2 \bar{\psi}_{xxxx} \varphi dx dt \right) + C \lambda \iint_{\tilde{\omega}_T} \xi \theta^2 |\varphi|^2 dx dt \\
(2.23) \quad &+ \delta \left(\iint_{Q_T} [\lambda^7 \mu^8 \xi^7 \theta^2 |\psi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\psi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\psi_{xx}|^2 + \lambda \mu^2 \xi \theta^2 |\psi_{xxx}|^2] dx dt \right).
\end{aligned}$$

Finally, for Ψ_4 we get

$$\begin{aligned}
\Psi_4 &= \operatorname{Re} \iint_{\tilde{\omega}_T} \eta \theta^2 \bar{\psi} (-g^0) dx dt \\
(2.24) \quad &\leq \delta \lambda^7 \mu^8 \iint_{Q_T} \xi^7 \theta^2 |\psi|^2 dx dt + C \lambda^{-7} \mu^{-8} \iint_{Q_T} \xi^{-7} \theta^2 |g^0|^2 dx dt.
\end{aligned}$$

Combining (2.10), (2.13), (2.17), (2.23) and (2.24) we get

$$\begin{aligned}
&\iint_{\mathcal{O}_T} \theta^2 |\psi|^2 dx dt \\
(2.25) \quad &\leq C \left(\lambda \iint_{\tilde{\omega}_T} \xi \theta^2 |\varphi|^2 dx dt + \lambda^{-3} \mu^{-4} \iint_{\tilde{\omega}_T} \xi^{-3} \theta^2 |\varphi|^2 dx dt + \lambda^{-5} \mu^{-8} \iint_{\tilde{\omega}_T} \xi^{-5} |\varphi|^2 dx dt \right) \\
&+ \operatorname{Re} \left(\iint_{\tilde{\omega}_T} \eta \theta^2 \varphi (i \bar{\psi}_t + \bar{\psi}_{xx} - \bar{\psi}_{xxxx}) dx dt \right) + C \lambda^{-7} \mu^{-8} \iint_{Q_T} \xi^{-7} \theta^2 |g^0|^2 dx dt \\
&+ \delta \left(\iint_{Q_T} [\lambda^7 \mu^8 \xi^7 \theta^2 |\psi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\psi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\psi_{xx}|^2 + \lambda \mu^2 \xi \theta^2 |\psi_{xxx}|^2] dx dt \right).
\end{aligned}$$

Since we get $i \bar{\psi}_t + \bar{\psi}_{xx} - \bar{\psi}_{xxxx} = \overline{-i \psi_t + \psi_{xx} - \psi_{xxxx}} = \bar{g}^1$ in Q_T and

$$(2.26) \quad \operatorname{Re} \left(\iint_{\tilde{\omega}_T} \eta \theta^2 \varphi \bar{g}^1 dx dt \right) \leq \delta \lambda^7 \mu^8 \iint_{Q_T} \xi^7 \theta^2 |\varphi|^2 dx dt + C \lambda^{-7} \mu^{-8} \iint_{Q_T} \xi^{-7} \theta^2 |g^1|^2 dx dt,$$

then, putting together (2.25) and (2.26), yields that

$$\begin{aligned}
&\iint_{\mathcal{O}_T} \theta^2 |\psi|^2 dx dt \leq C \lambda^{-7} \mu^{-8} \iint_{Q_T} \xi^{-7} \theta^2 (|g^0|^2 + |g^1|^2) dx dt + \delta \lambda^7 \mu^8 \iint_{Q_T} \xi^7 \theta^2 |\varphi|^2 dx dt \\
(2.27) \quad &+ C \left(\lambda \iint_{\tilde{\omega}_T} \xi \theta^2 |\varphi|^2 dx dt + \lambda^{-3} \mu^{-4} \iint_{\tilde{\omega}_T} \xi^{-3} \theta^2 |\varphi|^2 dx dt + \lambda^{-5} \mu^{-8} \iint_{\tilde{\omega}_T} \xi^{-5} |\varphi|^2 dx dt \right) \\
&+ \delta \left(\iint_{Q_T} [\lambda^7 \mu^8 \xi^7 \theta^2 |\psi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\psi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\psi_{xx}|^2 + \lambda \mu^2 \xi \theta^2 |\psi_{xxx}|^2] dx dt \right).
\end{aligned}$$

Combining (2.9) with (2.27) we obtain the following

$$\begin{aligned}
 & \iint_{Q_T} (\lambda^7 \mu^8 \xi^7 \theta^2 |\varphi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\varphi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\varphi_{xx}|^2 + \lambda \mu^2 \xi \theta^2 |\varphi_{xxx}|^2) dx dt \\
 & + \iint_{Q_T} (\lambda^7 \mu^8 \xi^7 \theta^2 |\psi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\psi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\psi_{xx}|^2 + \lambda \mu^2 \xi \theta^2 |\psi_{xxx}|^2) dx dt \\
 & \leq C \left(\iint_{Q_T} \theta^2 (|g^0|^2 + |g^1|^2) dx dt + \lambda^{-7} \mu^{-8} \iint_{Q_T} \xi^{-7} \theta^2 (|g^0|^2 + |g^1|^2) dx dt \right) \\
 & + C \left(\lambda \iint_{\tilde{\omega}_T} \xi \theta^2 |\varphi|^2 dx dt + \lambda^{-3} \mu^{-4} \iint_{\tilde{\omega}_T} \xi^{-3} \theta^2 |\varphi|^2 dx dt + \lambda^{-5} \mu^{-8} \iint_{\tilde{\omega}_T} \xi^{-5} |\varphi|^2 dx dt \right) \\
 & + \delta \left(\iint_{Q_T} [\lambda^7 \mu^8 \xi^7 \theta^2 (|\varphi|^2 + |\psi|^2) + \lambda^5 \mu^6 \xi^5 \theta^2 |\psi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\psi_{xx}|^2 + \lambda \mu^2 \xi \theta^2 |\psi_{xxx}|^2] dx dt \right) \\
 & + C \left(\lambda^3 \mu^3 \int_0^T (\xi^3 \theta^2 (|\varphi_{xx}|^2 + |\psi_{xx}|^2)) (t, L) dt + \lambda \mu \int_0^T (\xi \theta^2 (|\varphi_{xxx}|^2 + |\psi_{xxx}|^2)) (t, L) dt \right).
 \end{aligned}$$

Then, for λ, μ large enough and δ small enough we get

$$\begin{aligned}
 & \iint_{Q_T} (\lambda^7 \mu^8 \xi^7 \theta^2 |\varphi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\varphi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\varphi_{xx}|^2 + \lambda \mu^2 \xi \theta^2 |\varphi_{xxx}|^2) dx dt \\
 & + \iint_{Q_T} (\lambda^7 \mu^8 \xi^7 \theta^2 |\psi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\psi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\psi_{xx}|^2 + \lambda \mu^2 \xi \theta^2 |\psi_{xxx}|^2) dx dt \\
 (2.28) \quad & \leq C \left(\iint_{Q_T} \theta^2 (|g^0|^2 + |g^1|^2) dx dt + \lambda \iint_{\tilde{\omega}_T} \xi \theta^2 |\varphi|^2 dx dt \right) \\
 & + C \lambda^3 \mu^3 \int_0^T (\xi^3 \theta^2 (|\varphi_{xx}|^2 + |\psi_{xx}|^2)) (t, L) dt + C \lambda \mu \int_0^T (\xi \theta^2 (|\varphi_{xxx}|^2 + |\psi_{xxx}|^2)) (t, L) dt \\
 & =: \mathcal{I} + \mathcal{B}_1 + \mathcal{B}_2.
 \end{aligned}$$

Step 3: Estimates of the boundary terms.

Now, we will find an estimate for the boundary term on the right-hand side of (2.28), precisely, \mathcal{B}_1 and \mathcal{B}_2 . Using trace Theorem (note that $5/2 < 7/2$) we have

$$\begin{aligned}
 \mathcal{B}_1 &= C \lambda^3 \mu^3 \int_0^T (\xi^3 \theta^2 (|\varphi_{xx}|^2 + |\psi_{xx}|^2)) (t, L) dt \\
 (2.29) \quad & \leq C \lambda^3 \mu^3 \int_0^T (\xi^3 \theta^2) (t, L) \left(\|\varphi\|_{H^{\frac{5}{2}}(\tilde{\omega}_T)}^2 + \|\psi\|_{H^{\frac{5}{2}}(\tilde{\omega}_T)}^2 \right) dt \\
 & \leq C \lambda^3 \mu^3 \int_0^T (\xi^3 \theta^2) (t, L) \left(\|\varphi\|_{H^{\frac{7}{2}}(\tilde{\omega}_T)}^2 + \|\psi\|_{H^{\frac{7}{2}}(\tilde{\omega}_T)}^2 \right) dt
 \end{aligned}$$

and

$$\begin{aligned}
 \mathcal{B}_2 &= C \lambda \mu \int_0^T (\xi \theta^2 (|\varphi_{xxx}|^2 + |\psi_{xxx}|^2)) (t, L) dt \\
 (2.30) \quad & \leq C \lambda^3 \mu^3 \int_0^T (\xi^3 \theta^2) (t, L) \left(\|\varphi\|_{H^{\frac{7}{2}}(\tilde{\omega}_T)}^2 + \|\psi\|_{H^{\frac{7}{2}}(\tilde{\omega}_T)}^2 \right) dt.
 \end{aligned}$$

So, putting together (2.29) and (2.30), yields that

$$(2.31) \quad \mathcal{B}_1 + \mathcal{B}_2 \leq C \lambda^3 \mu^3 \int_0^T (\xi^3 \theta^2) (t, L) \left(\|\varphi\|_{H^{\frac{7}{2}}(\tilde{\omega}_T)}^2 + \|\psi\|_{H^{\frac{7}{2}}(\tilde{\omega}_T)}^2 \right) dt.$$

Using interpolation in the Sobolev spaces $H^s(\Omega)$, for $s \geq 0$, yields that

$$\begin{aligned}
\mathcal{B}_1 + \mathcal{B}_2 &\leq C_1 \lambda^3 \mu^3 \int_0^T (\xi^3 \theta^2)(t, L) \|\varphi\|_{H^{\frac{11}{3}}(\tilde{\omega}_T)}^{\frac{21}{11}} \|\varphi\|_{L^2(\tilde{\omega}_T)}^{\frac{1}{11}} dt \\
&\quad + C_1 \lambda^3 \mu^3 \int_0^T (\xi^3 \theta^2)(t, L) \|\psi\|_{H^{\frac{11}{3}}(\tilde{\omega}_T)}^{\frac{21}{11}} \|\psi\|_{L^2(\tilde{\omega}_T)}^{\frac{1}{11}} dt \\
&= C_1 \lambda^3 \mu^3 \int_0^T \left(\xi^{\frac{255}{22}} \xi^{-\frac{189}{22}} \theta^{\frac{86}{22}} \theta^{-\frac{42}{22}} \right)(t, L) \|\varphi\|_{H^{\frac{11}{3}}(\tilde{\omega}_T)}^{\frac{21}{11}} \|\varphi\|_{L^2(\tilde{\omega}_T)}^{\frac{1}{11}} dt \\
&\quad + C_1 \lambda^3 \mu^3 \int_0^T \left(\xi^{\frac{255}{22}} \xi^{-\frac{189}{22}} \theta^{\frac{86}{22}} \theta^{-\frac{42}{22}} \right)(t, L) \|\psi\|_{H^{\frac{11}{3}}(\tilde{\omega}_T)}^{\frac{21}{11}} \|\psi\|_{L^2(\tilde{\omega}_T)}^{\frac{1}{11}} dt \\
&\leq C_\epsilon \lambda^6 \mu^6 \int_0^T (\xi^{255} \theta^{86})(t, L) \|\varphi\|_{L^2(\tilde{\omega}_T)}^2 dt + \epsilon \lambda^{-2} \mu^{-2} \int_0^T \left(\xi^{-\frac{189}{21}} \theta^{-2} \right)(t, L) \|\varphi\|_{H^{\frac{11}{3}}(\tilde{\omega}_T)}^2 dt \\
&\quad + C_\epsilon \lambda^6 \mu^6 \int_0^T (\xi^{255} \theta^{86})(t, L) \|\psi\|_{L^2(\tilde{\omega}_T)}^2 dt + \epsilon \lambda^{-2} \mu^{-2} \int_0^T \left(\xi^{-\frac{189}{21}} \theta^{-2} \right)(t, L) \|\psi\|_{H^{\frac{11}{3}}(\tilde{\omega}_T)}^2 dt,
\end{aligned}$$

or equivalently,

$$\begin{aligned}
\mathcal{B}_1 + \mathcal{B}_2 &\leq C_\epsilon \lambda^6 \mu^6 \int_0^T (\xi^{255} \theta^{86})(t, L) \|\varphi\|_{L^2(\tilde{\omega}_T)}^2 dt \\
&\quad + \epsilon \lambda^{-2} \mu^{-2} \int_0^T (\xi^{-9} \theta^{-2})(t, L) \|\varphi\|_{H^{\frac{11}{3}}(\tilde{\omega}_T)}^2 dt \\
(2.32) \quad &\quad + C_\epsilon \lambda^6 \mu^6 \int_0^T (\xi^{255} \theta^{86})(t, L) \|\psi\|_{L^2(\tilde{\omega}_T)}^2 dt \\
&\quad + \epsilon \lambda^{-2} \mu^{-2} \int_0^T (\xi^{-9} \theta^{-2})(t, L) \|\psi\|_{H^{\frac{11}{3}}(\tilde{\omega}_T)}^2 dt \\
&=: \mathcal{I}_1 + \mathcal{I}_2 + \mathcal{I}_3 + \mathcal{I}_4,
\end{aligned}$$

for some positive constant C_ϵ .

At this moment, our goal is to prove integrals \mathcal{I}_i , for $i = 1, 2, 3$, can be absolved by the left-hand side of (2.28). Let us start with the analysis of \mathcal{I}_2 , precisely the quantity

$$\int_0^T (\xi^{-9} \theta^{-2})(t, L) \|\varphi\|_{H^{\frac{11}{3}}(\tilde{\omega}_T)}^2 dt.$$

Consider $\varphi_1(t, x) := \xi_1(t) \varphi(t, x)$ with

$$\xi_1(t) = \theta^{-1} \xi^{-\frac{1}{2}}.$$

Then φ_1 satisfies the system

$$(2.33) \quad \begin{cases} -i\varphi_{1t} + \varphi_{1xx} - \varphi_{1xxxx} = f_1 := \xi_{1t}\varphi, & \text{in } Q_T, \\ \varphi_1(t, 0) = \varphi_1(t, L) = \varphi_{1x}(t, 0) = \varphi_{1x}(t, L) = 0, & \text{on } (0, T), \\ \varphi_1(T, x) = 0, & \text{in } \Omega. \end{cases}$$

Now, observe that, since $\varphi_x(t, 0) = 0$ and $|\xi_{1t}| \leq C \lambda \xi^{\frac{3}{2}} \theta^{-1}$, we have

$$\begin{aligned}
\|f_1\|_{L^2(Q_T)}^2 &\leq C \iint_{Q_T} \theta^{-2} \lambda^2 \xi^3 |\varphi|^2 dx dt \\
(2.34) \quad &\leq C \iint_{Q_T} \{ \lambda^2 \xi^3 |\varphi|^2 + \lambda^3 |\varphi_x|^2 + \lambda |\varphi_{xx}|^2 + \lambda^{-1} |\varphi_{xxx}|^2 \} \theta^{-2} dx dt,
\end{aligned}$$

for some constant $C > 0$ and all $s \geq s_0$. Moreover, thanks to Section A, $\varphi_1 \in L^2(0, T; H^2(\Omega)) \cap C([0, T]; L^2(\Omega))$. Then, interpolating between $L^2(0, T; H^2(\Omega))$ and $L^\infty(0, T; L^2(\Omega))$, we infer that $\varphi_1 \in L^2(0, T; H^{5/3}(\Omega))$ and

$$(2.35) \quad \|\varphi_1\|_{L^2(0, T; H^{5/3}(\Omega))} \leq C \|f_1\|_{L^2(Q_T)}.$$

Let $\varphi_2(t, x) := \xi_2(t)\varphi(t, x)$ with

$$\xi_2 = \theta^{-1}\xi^{-\frac{5}{2}}.$$

Then v_2 satisfies system (2.33) with f_1 replaced by

$$f_2 := \xi_{2t}\xi_1^{-1}\varphi_1.$$

Observe that $|\theta_{2t}\theta_1^{-1}| \leq C\lambda$. Thus, we obtain

$$(2.36) \quad \|f_2\|_{L^2(0,T;H^{5/3}(\Omega))} \leq C\lambda \|\varphi_1\|_{L^2(0,T;H^{5/2}(\Omega))}.$$

Now, by using that φ_2 belongs to $L^2(0, T; H^4(\Omega))$ and $L^\infty(0, T; H^2(\Omega))$, thanks to (A.4), and interpolating these two spaces, we have that

$$\varphi_2 \in L^2(0, T; H^{11/3}(\Omega)) \cap L^\infty(0, T; H^{8/3}(\Omega))$$

with

$$(2.37) \quad \|\varphi_2\|_{L^2(0,T;H^{11/3}(\Omega)) \cap L^\infty(0,T;H^{8/3}(\Omega))} \leq C \|f_2\|_{L^2(0,T;H^{5/3}(\Omega))}.$$

Thus we infer from (2.35)–(2.37), the following

$$(2.38) \quad \begin{aligned} \|\varphi_2\|_{L^2(0,T;H^{11/3}(\Omega))}^2 &\leq C_1\lambda \|f_1\|_{L^2(Q_T)}^2 \\ &\leq C_2 \int_{Q_T} (\lambda^3 \xi^3 |\varphi|^2 + \lambda^4 |\varphi_x|^2 + \lambda^2 |\varphi_{xx}|^2 + |\varphi_{xxx}|^2) \theta^{-2} dx dt. \end{aligned}$$

Hence, replacing $\varphi_2 = \theta^{-1}\varphi^{-\frac{9}{2}}$ in (2.38), for some constant $C_3 > 0$, yields that

$$(2.39) \quad \begin{aligned} \int_0^T (\xi^{-9}\theta^{-2})(t, L) \|\varphi(t, \cdot)\|_{H^{11/3}(\tilde{\omega}_T)}^2 dt \\ \leq C_3 \int_{Q_T} (\lambda^3 \xi^3 |\varphi|^2 + \lambda^4 |\varphi_x|^2 + \lambda^2 |\varphi_{xx}|^2 + |\varphi_{xxx}|^2) \theta^{-2} dx dt. \end{aligned}$$

Note that analogously we can infer the same relation for ψ , that is,

$$(2.40) \quad \begin{aligned} \int_0^T (\xi^{-9}\theta^{-2})(t, L) \|\psi(t, \cdot)\|_{H^{11/3}(\tilde{\omega}_T)}^2 dt \\ \leq C_3 \int_{Q_T} (\lambda^3 \xi^3 |\psi|^2 + \lambda^4 |\psi_x|^2 + \lambda^2 |\psi_{xx}|^2 + |\psi_{xxx}|^2) \theta^{-2} dx dt. \end{aligned}$$

Therefore, adding (2.39) and (2.40), putting in (2.32) and, finally, comparing with (2.28), for λ and μ large enough, yields that,

$$(2.41) \quad \begin{aligned} &\iint_{Q_T} (\lambda^7 \mu^8 \xi^7 \theta^2 |\varphi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\varphi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\varphi_{xx}|^2 + \lambda \mu^2 \xi \theta^2 |\varphi_{xxx}|^2) dx dt \\ &+ \iint_{Q_T} (\lambda^7 \mu^8 \xi^7 \theta^2 |\psi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\psi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\psi_{xx}|^2 + \lambda \mu^2 \xi \theta^2 |\psi_{xxx}|^2) dx dt \\ &\leq C \left(\iint_{Q_T} \theta^2 (|g^0|^2 + |g^1|^2) dx dt + \lambda \iint_{\tilde{\omega}_T} \xi \theta^2 |\varphi|^2 dx dt \right), \end{aligned}$$

since $|(\xi^{255}\theta^{86})(t, L)|$ is bounded in terms of $t \in [0, T]$ due to the choices in (2.1), what guarantees (2.3), and so the Carleman is shown. \square

3. NULL CONTROLLABILITY RESULTS

In this section we prove the existence of insensitizing controls for the linearized system (1.6). First we need to obtain an estimate as in Theorem 2.2, with weights that remain bounded as $t \rightarrow T$, i.e., have blow-up only in $t = 0$. To this purpose we introduce the new weights

$$(3.1) \quad \begin{aligned} \sigma &= e^m, & \nu(t, x) &= \frac{e^{3\mu\eta(x)}}{\gamma(t)} & \text{and} & & m(t, x) &= \lambda \frac{e^{3\mu\eta(x)} - e^{5\mu\|\eta\|_\infty}}{\gamma(t)}, \\ \sigma^* &= e^{m^*}, & \nu^*(t) &= \min_{x \in \bar{\Omega}} \nu(x, t) & \text{and} & & m^*(t) &= \min_{x \in \bar{\Omega}} \nu(x, t), \\ \hat{\sigma} &= e^{\hat{m}}, & \hat{\nu}(t) &= \max_{x \in \bar{\Omega}} \nu(x, t) & \text{and} & & \hat{m}(t) &= \max_{x \in \bar{\Omega}} \nu(x, t), \end{aligned}$$

where γ is given by

$$(3.2) \quad \gamma(t) = \begin{cases} t(T-t), & 0 \leq t \leq T/2, \\ T^2/4, & T/2 < t \leq T. \end{cases}$$

Combining Carleman estimate (2.3) with classical energy estimates for the fourth order Schrödinger system, satisfied by φ and ψ , we can prove the following result.

Proposition 3.1. *With the hypothesis of Proposition 2.2 the solution (φ, ψ) of (2.4) satisfies the following*

$$\begin{aligned} & \|\varphi\|_{L^2(T/2, T; L^2(\Omega))} + \|\varphi_x\|_{L^2(T/2, T; L^2(\Omega))} + \|\varphi_{xx}\|_{L^2(T/2, T; L^2(\Omega))} \\ & \leq \|(\varphi, \psi)\|_{(L^2(T/4, T/2; L^2(\Omega)))^2} + \|(g^0, g^1)\|_{(L^2(T/2, T; L^2(\Omega)))^2} \end{aligned}$$

and

$$\begin{aligned} & \|\psi\|_{L^2(T/2, T; L^2(\Omega))} + \|\psi_x\|_{L^2(T/2, T; L^2(\Omega))} + \|\psi_{xx}\|_{L^2(T/2, T; L^2(\Omega))} \\ & \leq \|\psi\|_{L^2(T/4, T/2; L^2(\Omega))} + \|g^1\|_{L^2(T/2, T; L^2(\Omega))}. \end{aligned}$$

Proof. Let us consider $\kappa \in C^1([0, T])$ such that

$$\kappa = \begin{cases} 0, & \text{if } t \in [0, T/4], \\ 1, & \text{if } t \in [T/2, T]. \end{cases}$$

Note that if (φ, ψ) is a solution for (2.4), so $(\kappa\varphi, \kappa\psi)$ satisfies the following system

$$(3.3) \quad \begin{cases} i(\kappa\varphi)_t + (\kappa\varphi)_{xx} - (\kappa\varphi)_{xxxx} = 1_{\mathcal{O}}(\kappa\psi) + \kappa g^0 + i\kappa_t\varphi, & \text{in } Q_T, \\ i(\kappa\psi)_t + (\kappa\psi)_{xx} - (\kappa\psi)_{xxxx} = \kappa g^1 + i\kappa_t\psi, & \text{in } Q_T, \\ (\kappa\varphi)(t, 0) = (\kappa\varphi)(t, L) = (\kappa\varphi)_x(t, 0) = (\kappa\varphi)_x(t, L) = 0, & \text{on } t \in (0, T), \\ \psi(t, 0) = (\kappa\psi)(t, L) = (\kappa\psi)_x(t, 0) = (\kappa\psi)_x(t, L) = 0, & \text{on } t \in (0, T), \\ (\kappa\varphi)(T, x) = 0, (\kappa\psi)(T, x) = 0, & \text{in } \Omega. \end{cases}$$

Now, since $\kappa, \kappa_t \in C([0, T])$ and $C([0, T]) \hookrightarrow L^\infty(0, T)$, moreover, $\kappa\psi, \kappa_t\psi \in L^2(0, T; H_0^2(\Omega))$. Then, for $g^1 \in L^2(0, T; H_0^2(\Omega))$ we get that $\kappa\psi$ satisfy a fourth order Schrödinger system equation with null data and right-hand side in $L^2(0, T; H_0^2(\Omega))$. Therefore, we get that

$$(3.4) \quad \begin{aligned} & \int_{\Omega} |\kappa(t)\psi(t)| dx + \iint_{Q_T} |\kappa\psi_x| dx dt + \iint_{Q_T} |\kappa\psi_{xx}| dx dt \\ & \leq C \left(\iint_{Q_T} |\kappa g^1| dx dt + \iint_{Q_T} |\kappa_t\psi| dx dt \right). \end{aligned}$$

Multiplying the first equation of (3.3) by $\kappa\varphi$ and integrating over Ω we obtain, after taking the real part and using Young inequality for the integral term of $1_{\mathcal{O}}\kappa\psi\kappa\varphi$, that

$$(3.5) \quad \begin{aligned} & -\frac{1}{2} \frac{d}{dt} \int_{\Omega} |\kappa(t)\varphi(t)|^2 + \int_{\Omega} |\kappa\varphi_x|^2 dx + \int_{\Omega} |\kappa\varphi_{xx}|^2 dx \\ & \leq C \left(\int_{\Omega} |\kappa g^0|^2 dx + \int_{\Omega} |\kappa_t\varphi|^2 dx + \int_{\Omega} |\kappa\psi|^2 dx \right) + \delta \int_{\mathcal{O}} |\kappa\varphi|^2 dx. \end{aligned}$$

Finally, integrating (3.5) in $[t, T]$, combining with (3.4) and taking δ small enough, we get

$$(3.6) \quad \begin{aligned} & \int_{\Omega} |\kappa(t)\varphi(t)|^2 dx + \iint_{Q_T} |\kappa\varphi_x|^2 dxdt + \iint_{Q_T} |\kappa\varphi_{xx}|^2 dxdt \\ & \leq C \left(\iint_{Q_T} |\kappa|^2 (|g^0|^2 + |g^1|^2) dxdt + \iint_{Q_T} |\kappa_t|^2 (\varphi^2 + |\psi|^2) dxdt \right). \end{aligned}$$

Therefore, Proposition 3.1 is a consequence of equations (3.4) and (3.6). \square

As a consequence of the previous result, and due to the definition of (3.1), the following Carleman estimate, with new weight functions σ and ν , can be obtained.

Proposition 3.2. *There exists a constant $C(s, \lambda) := C > 0$, such that every solution (φ, ψ) of (2.4) satisfies*

$$(3.7) \quad \begin{aligned} & \iint_{Q_T} (\lambda^7 \mu^8 \nu^7 \sigma^2 |\varphi|^2 + \lambda^5 \mu^6 \nu^5 \sigma^2 |\varphi_x|^2 + \lambda^3 \mu^4 \nu^3 \sigma^2 |\varphi_{xx}|^2) dxdt \\ & + \iint_{Q_T} (\lambda^7 \mu^8 \nu^7 \sigma^2 |\psi|^2 + \lambda^5 \mu^6 \nu^5 \sigma^2 |\psi_x|^2 + \lambda^3 \mu^4 \nu^3 \sigma^2 |\psi_{xx}|^2) dxdt \\ & \leq C \left(\iint_{Q_T} \sigma^2 (|g^0|^2 + |g^1|^2) dxdt + \lambda \iint_{\tilde{\omega}_T} \nu \sigma^2 |\varphi|^2 dxdt \right). \end{aligned}$$

Proof. The result is consequence of Proposition 3.1. Indeed, noting that $\xi = \nu$ and $l = m$, for $t \in [0, T/2]$, and since l is constant in $[T/2, T]$, yields that

$$(3.8) \quad \begin{aligned} & \int_0^{T/2} \int_{\Omega} (\lambda^7 \mu^8 \nu^7 \sigma^2 |\varphi|^2 + \lambda^5 \mu^6 \nu^5 \sigma^2 |\varphi_x|^2 + \lambda^3 \mu^4 \nu^3 \sigma^2 |\varphi_{xx}|^2) dxdt \\ & + \int_0^{T/2} \int_{\Omega} (\lambda^7 \mu^8 \nu^7 \sigma^2 |\psi|^2 + \lambda^5 \mu^6 \nu^5 \sigma^2 |\psi_x|^2 + \lambda^3 \mu^4 \nu^3 \sigma^2 |\psi_{xx}|^2) dxdt \\ & = \int_0^{T/2} \int_{\Omega} (\lambda^7 \mu^8 \xi^7 \theta^2 |\varphi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\varphi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\varphi_{xx}|^2) dxdt \\ & + \int_0^{T/2} \int_{\Omega} (\lambda^7 \mu^8 \xi^7 \theta^2 |\psi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\psi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\psi_{xx}|^2) dxdt. \end{aligned}$$

Additionally, for $t \in [T/2, T]$, we have that

$$(3.9) \quad \begin{aligned} & \int_{T/2}^T \int_{\Omega} (\lambda^7 \mu^8 \xi^7 \theta^2 |\varphi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\varphi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\varphi_{xx}|^2) dxdt \\ & + \int_{T/2}^T \int_{\Omega} (\lambda^7 \mu^8 \xi^7 \theta^2 |\psi|^2 + \lambda^5 \mu^6 \xi^5 \theta^2 |\psi_x|^2 + \lambda^3 \mu^4 \xi^3 \theta^2 |\psi_{xx}|^2) dxdt \\ & \leq C \left(\int_{T/2}^T \int_{\Omega} (|\varphi|^2 + |\varphi_x|^2 + |\varphi_{xx}|^2 + |\psi|^2 + |\psi_x|^2 + |\psi_{xx}|^2) dxdt \right) \\ & \leq C \left(\int_{T/4}^{T/2} \int_{\Omega} (|\varphi|^2 + |\psi|^2) dxdt + \int_{T/2}^T \int_{\Omega} (|g^0|^2 + |g^1|^2) dxdt \right) \\ & \leq C \left(\int_{T/4}^{T/2} \int_{\Omega} \xi^7 \theta^2 (|\varphi|^2 + |\psi|^2) dxdt + \int_{T/2}^T \int_{\Omega} \sigma^2 (|g^0|^2 + |g^1|^2) dxdt \right), \end{aligned}$$

thanks to Proposition 3.1.

Finally, we note that

$$(3.10) \quad \begin{aligned} & \iint_{Q_T} \theta^2 (|g^0|^2 + |g^1|^2) dx dt + \lambda \iint_{\tilde{\omega}_T} \xi \theta^2 |\varphi|^2 dx dt \\ & \leq C \left(\iint_{Q_T} \sigma^2 (|g^0|^2 + |g^1|^2) dx dt + \lambda \iint_{\tilde{\omega}_T} \nu \sigma^2 |\varphi|^2 dx dt \right). \end{aligned}$$

Thus, the result follows from (2.3), (3.8), (3.9) and (3.10). \square

Remark 3.1. We point out that Proposition 3.1 holds true by taking the minimum of the weights in the left-hand side and maximum of the weights in the right-hand side of (3.4).

3.1. Null controllability: Linear case. In what follows we use (3.7), from Proposition 3.2, to deduce the desired null controllability property. Denote $\mathcal{L} = \mathcal{L}^* = i\partial_t + \partial_{xx} - \partial_{xxxx}$ and introduce the following space

$$\begin{aligned} \mathcal{C} = \Big\{ & (u, v, h); (\hat{\sigma})^{-1}u \in L^2(Q_T), (\hat{\sigma})^{-1}v \in L^2(0, T; H^{-2}(\Omega)), (\hat{\nu})^{-\frac{1}{2}}(\hat{\sigma})^{-1}h \in L^2(Q_T), \\ & (\nu^*)^{-\frac{7}{2}}(\sigma^*)^{-1}(\mathcal{L}u - 1_\omega h) \in L^2(Q_T), (\nu^*)^{-\frac{7}{2}}(\sigma^*)^{-1}(\mathcal{L}^*v - 1_\mathcal{O}u) \in L^2(0, T; H^{-2}(\Omega)), \\ & (\hat{\nu})^{-2}\hat{\sigma}^{-1}u \in L^2(0, T; H^4(\Omega)) \cap L^\infty(0, T; H_0^2(\Omega)), \\ & (\hat{\nu})^{-2}\hat{\sigma}^{-1}v \in L^2(0, T; H_0^2(\Omega)) \cap L^\infty(0, T; L^2(\Omega)), v|_{t=T} = 0 \text{ in } \Omega \Big\}. \end{aligned}$$

Remark 3.2. It is important to observe that \mathcal{C} is a Banach space endowed with its natural norm. Additionally, as consequence from the definition of the set \mathcal{C} , an element $(u, v, h) \in \mathcal{C}$ is such that $v|_{t=0} = 0$ in Ω . This holds since $(\hat{\nu})^{-2}\hat{\sigma}^{-1}v$ belongs to $L^\infty(0, T; L^2(\Omega))$ and $(\hat{\nu})^{-2}\hat{\sigma}^{-1}$ blow-up only at $t = 0$.

We are now in position to prove the null controllability property for solutions of (1.6). The result can be read as follows.

Theorem 3.3. Assume the same hypothesis of Proposition 3.2. Additionally, consider

$$(\nu^*)^{-\frac{7}{2}}(\hat{\sigma})^{-1}f^0 \in L^2(Q_T) \quad \text{and} \quad (\nu^*)^{-\frac{7}{2}}(\hat{\sigma})^{-1}f^1 \in L^2(0, T; H^{-2}(\Omega)).$$

Therefore, we can find a control $h(x, t) = h$ such that the associated solution (u, v) of

$$(3.11) \quad \begin{cases} iu_t + u_{xx} - u_{xxxx} = f^0 + 1_\omega h, & \text{in } Q_T, \\ iv_t + v_{xx} - v_{xxxx} = f^1 + 1_\mathcal{O}u, & \text{in } Q_T, \\ u(t, 0) = u(t, L) = v(t, 0) = v(t, L) = 0, & \text{on } t \in (0, T), \\ u_x(t, 0) = u_x(t, L) = v_x(t, 0) = v_x(t, L) = 0, & \text{on } t \in (0, T), \\ u(0, x) = u_0(x), v(T, x) = 0, & \text{in } \Omega, \end{cases}$$

satisfies $(u, v, h) \in \mathcal{C}$. In particular, $v|_{t=0} \equiv 0$ in Ω .

Proof. We introduce the following spaces

$$R_0 = \{u \in H_0^2(\Omega); u_{xxxx} \in H_0^2(\Omega)\},$$

$$Y_0 = C([0, T]; H^4(\Omega)) \cap C^1([0, T]; H_0^2(\Omega))$$

and

$$Y_1 = C([0, T]; H_0^2(\Omega)) \cap C^1([0, T]; H^{-2}(\Omega)).$$

Also, let us consider

$$P_0 = \{(\varphi, \psi) \in Y_1 \times Y_0 : \mathcal{L}^*v - 1_\mathcal{O}u \in L^2(Q_T)\}.$$

Thanks to Theorem A.2, P_0 is nonempty. Moreover, from now on we will use \mathcal{L} instead of \mathcal{L}^* , since both are equal.

Now, define the bilinear form $a : P_0 \times P_0 \rightarrow \mathbb{R}$ by

$$a((\tilde{\varphi}, \tilde{\psi}), (\varphi, \psi)) := \operatorname{Re} \left(\iint_{Q_T} (\hat{\sigma})^2 (\mathcal{L}\hat{\varphi} - 1_{\mathcal{O}}\hat{\psi}) (\overline{\mathcal{L}\varphi - 1_{\mathcal{O}}\psi}) dxdt + \iint_{Q_T} (\hat{\sigma})^2 (\mathcal{L}\hat{\psi}) (\overline{\mathcal{L}\varphi}) dxdt \right) \\ + \operatorname{Re} \left(\iint_{\tilde{\omega}_T} \hat{\nu}(\hat{\sigma})^2 \hat{\varphi} \overline{\varphi} dxdt \right),$$

and the linear form $G : P_0 \rightarrow \mathbb{R}$ given by

$$\langle G, (\varphi, \psi) \rangle := \operatorname{Re} \iint_{Q_T} f^0 \overline{\varphi} dxdt + \int_0^T \langle f^1, \overline{\psi} \rangle dt,$$

where $\langle \cdot, \cdot \rangle$ denotes the duality between $H^{-2}(\Omega)$ and $H_0^2(\Omega)$. Thanks to Proposition 3.2, the bilinear form over $P_0 \times P_0$, is sesquilinear, positive and coercive. Let P be the completion of P_0 with the norm induced by $a(\cdot, \cdot)$, in this case, $a(\cdot, \cdot)$ is well-defined, continuous and coercive on $P \times P$. Now, note that for all $(\varphi, \psi) \in P_0$ we have

$$\begin{aligned} \langle G, (\varphi, \psi) \rangle &= \operatorname{Re} \iint_{Q_T} f^0 \overline{\varphi} dxdt + \int_0^T \langle f^1, \overline{\psi} \rangle dt \\ &\leq \left(\iint_{Q_T} (\nu^*)^7 (\sigma^*)^2 (|\varphi|^2 + |\psi|^2) dxdt \right)^{1/2} \\ &\quad \times \left(\iint_{Q_T} (\nu^*)^{-7} (\sigma^*)^{-2} |f^0|^2 dxdt + \int_0^T (\nu^*)^{-7} (\sigma^*)^{-2} \|f^1\|_{H^{-2}(\Omega)}^2 dt \right)^{1/2} \\ &\leq C a((\varphi, \psi), (\varphi, \psi))^{1/2} \left(\iint_{Q_T} (\nu^*)^{-7} (\sigma^*)^{-2} |f^0|^2 dxdt + \int_0^T (\nu^*)^{-7} (\sigma^*)^{-2} \|f^1\|_{H^{-2}(\Omega)}^2 dt \right)^{1/2}, \end{aligned}$$

where we use Young inequality on the first inequality. Therefore, G is a bounded functional on P_0 and we can extend it continuously to a bounded functional on P due to the Hahn-Banach theorem. Hence, from Lax-Milgram's lemma, we deduce that the variational problem

$$(3.12) \quad \begin{cases} \text{Find } (\hat{\varphi}, \hat{\psi}) \in P \text{ such that, } \forall (\varphi, \psi) \in P, \\ a((\hat{\varphi}, \hat{\psi}), (\varphi, \psi)) = \langle G, (\varphi, \psi) \rangle, \end{cases}$$

has a unique solution $(\hat{\varphi}, \hat{\psi}) \in P \times P$.

Let us define $(\hat{u}, \hat{v}, \hat{h})$ by

$$(3.13) \quad \begin{cases} \hat{u} = (\hat{\sigma})^2 (\mathcal{L}\hat{\varphi} - 1_{\mathcal{O}}\hat{\psi}), & \text{in } Q_T, \\ \hat{v} = (\hat{\sigma})^2 \mathcal{L}\hat{\psi}, & \text{in } Q_T, \\ \hat{h} = -\hat{\nu}(\hat{\sigma})^2 \hat{\varphi}, & \text{in } Q_T, \end{cases}$$

remembering that $\mathcal{L}^* = \mathcal{L}$. Thanks to (3.12) and (3.13), we have that

$$\iint_{Q_T} (\hat{\sigma})^{-2} (|\hat{u}|^2 + |\hat{v}|^2 + (\hat{\nu})^{-1} |\hat{h}|^2) dxdt = a((\hat{\varphi}, \hat{\psi}), (\hat{\varphi}, \hat{\psi})) < \infty.$$

Considering (\tilde{u}, \tilde{v}) be a weak solution of

$$(3.14) \quad \begin{cases} i\tilde{u}_t + \tilde{u}_{xx} - \tilde{u}_{xxx} = f^0 + 1_{\omega}\tilde{h}, & \text{in } Q_T, \\ i\tilde{v}_t + \tilde{v}_{xx} - \tilde{v}_{xxx} = f^1 + 1_{\mathcal{O}}\tilde{u}, & \text{in } Q_T, \\ \tilde{u}(t, 0) = \tilde{u}(t, L) = \tilde{v}(t, 0) = \tilde{v}(t, L) = 0, & \text{on } t \in (0, T), \\ \tilde{u}_x(t, 0) = \tilde{u}_x(t, L) = \tilde{v}_x(t, 0) = \tilde{v}_x(t, L) = 0, & \text{on } t \in (0, T), \\ \tilde{u}(0, x) = 0, \tilde{v}(T, x) = 0, & \text{in } \Omega, \end{cases}$$

with control $h = \hat{h}$ and source terms f^0 and f^1 , since $\tilde{h} \in L^2(Q_T)$, we have, from well-posed result given by Theorem A.2, that (\tilde{u}, \tilde{v}) are well defined. In the following we prove that the weak solution

(\hat{u}, \hat{u}) is a solution by transposition. In fact, for every $(\varphi, \psi) \in P_0$, it holds from (3.12) and (3.13) that

$$(3.15) \quad \begin{aligned} & \operatorname{Re} \iint_{Q_T} f^0 \bar{\varphi} dx dt + \int_0^T \langle f^1, \bar{\psi} \rangle_{H^{-2} \times H_0^2} dt + \operatorname{Re} \iint_{\tilde{\omega}} \hat{h} \bar{\varphi} dx dt \\ &= \operatorname{Re} \iint_{Q_T} \hat{u} (\overline{\mathcal{L}\varphi - 1_{\mathcal{O}}\psi}) dx dt + \int_0^T \langle \hat{v}, \overline{\mathcal{L}\psi} \rangle_{H^{-2} \times H_0^2} dt. \end{aligned}$$

From (3.15), we get that

$$\begin{aligned} \operatorname{Re} \iint_{Q_T} \hat{u} \bar{g}^0 dx dt + \int_0^T \langle \hat{v}, \bar{g}^1 \rangle_{H^{-2} \times H_0^2} dt &= \operatorname{Re} \iint_{\tilde{\omega}} \hat{h} \bar{\varphi} dx dt \\ &+ \operatorname{Re} \iint_{Q_T} f^0 \bar{\varphi} dx dt + \int_0^T \langle f^1, \bar{\psi} \rangle_{H^{-2} \times H_0^2} dt, \end{aligned}$$

for all $(g^0, g^1) \in L^2(0, T; H_0^1(\Omega))$, that is, $(\hat{u}, \hat{v}) = (\tilde{u}, \tilde{v})$.

Now on, we prove that solutions \hat{u} and \hat{v} of (3.14) are, in fact, more regular. Let us start defining the functions

$$\begin{aligned} u_* &:= (\hat{\nu})^{-2}(\hat{\sigma})^{-1}\hat{u}, & v_* &:= (\hat{\nu})^{-2}(\hat{\sigma})^{-1}\hat{v}, \\ f_*^0 &:= (\hat{\nu})^{-2}(\hat{\sigma})^{-1}(f^0 + \hat{h}1_{\omega}) & \text{and} & \quad f_*^1 := (\hat{\nu})^{-2}(\hat{\sigma})^{-1}f^1. \end{aligned}$$

It follows, from (3.11), that u_*, v_*, f_*^1 and f_*^2 satisfies the following system

$$(3.16) \quad \begin{cases} i(u_*)_t + (u_*)_{xx} - (u_*)_{xxxx} = f_*^0 + i((\hat{\nu})^{-2}(\hat{\sigma})^{-1})_t \hat{u}, & \text{in } Q_T, \\ i(v_*)_t + (v_*)_{xx} - (v_*)_{xxxx} = f_*^1 + 1_{\mathcal{O}}u_* + i((\hat{\nu})^{-2}(\hat{\sigma})^{-1})_t \hat{v}, & \text{in } Q_T, \\ u_* = v_* = 0, & \text{in } \Sigma, \\ (u_*)_{t=0} = 0, (v_*)_{t=T} = 0, & \text{in } \Omega. \end{cases}$$

Now, since $(\hat{\nu})^{-2}(\hat{\sigma})^{-1})_t \leq CT^2 s(\hat{\sigma})^{-1}$ we get that $f_*^0 + i((\hat{\nu})^{-2}(\hat{\sigma})^{-1})_t \hat{u} \in L^2(Q)$ and also $f_*^1 + i((\hat{\nu})^{-2}(\hat{\sigma})^{-1})_t \hat{v} \in L^2(0, T; H^{-2}(\Omega))$. Now, using the results of Section A, for (3.16), we obtain

$$u_* \in L^2(0, T; H^4(\Omega)) \cap L^\infty(0, T; H_0^2(\Omega))$$

and

$$v_* \in L^2(0, T; H^2(\Omega)) \cap L^\infty(0, T; L^2(\Omega)).$$

This finishes the proof of Theorem 3.3. \square

3.2. Null controllability: Nonlinear case. In this section we use an inverse mapping theorem to obtain the existence of insensitizing controls for the fourth order nonlinear Schrödinger equation (1.5). We invite the reader to see the result below as well as additional comments on [2].

Theorem 3.4 (Inverse Mapping Theorem). *Let B_1 and B_2 be two Banach spaces and let*

$$\mathcal{Y} : B_1 \rightarrow B_2$$

satisfying $\mathcal{Y} \in C^1(B_1, B_2)$. Assume that $b_1 \in B_1$, $\mathcal{Y}(b_1) = b_2$ and

$$\mathcal{Y}'(b_1) : B_1 \rightarrow B_2$$

is surjective. Then, there exists $\delta > 0$ such that, for every $b' \in B_2$ satisfying

$$\|b' - b_2\|_{B_2} < \delta,$$

there exists a solution of the equation

$$\mathcal{Y}(b) = b', \quad b \in B_1.$$

Finally, we will give the proof of the main result of this manuscript.

Proof of Theorem 1.2. Consider, in Theorem 3.4, the following

$$B_1 = \mathcal{C} \quad \text{and} \quad B_2 = L^2((\hat{\nu})^{-6}(\hat{\sigma})^{-3}(0, T); L^2(\Omega)) \times L^2((\hat{\nu})^{-6}(\hat{\sigma})^{-3}(0, T); H^{-2}(\Omega)).$$

Define the operator

$$\mathcal{Y} : B_1 \rightarrow B_2$$

such that

$$\mathcal{Y}(u, v, h) := (\mathcal{L}u - \zeta|u|^2u - 1_\omega h, \mathcal{L}v - \bar{\zeta}\bar{u}^2\bar{v} - \bar{\zeta}|u|^2v - 1_{\mathcal{O}}u).$$

Claim 1. *Operator \mathcal{Y} belongs to $C^1(B_1, B_2)$.*

Indeed, first note that all terms of \mathcal{Y} are linear except: $|u|^2u$, $\bar{u}^2\bar{v}$ and $|u|^2v$. So, the Claim 1 is equivalent to prove that the trilinear operator given by

$$((u_1, v_1, h_1), (u_1, v_1, h_1), (u_1, v_1, h_1)) \mapsto u_1 u_2 u_3$$

and

$$((u_1, v_1, h_1), (u_1, v_1, h_1), (u_1, v_1, h_1)) \mapsto u_1 u_2 v_3,$$

are continuous maps from \mathcal{C}^3 to $L^2((\hat{\nu})^{-6}(\hat{\sigma})^{-1}(0, T); L^2(\Omega))$. However, $(u_i, v_i, h_i) \in \mathcal{C}$, thus we get that

$$(\hat{\nu})^{-2}(\hat{\sigma})^{-1}u \in L^2(0, T; H^4((\Omega)) \cap L^\infty(0, T; H_0^2(\Omega))) \hookrightarrow L^6(Q_T)$$

and

$$(\hat{\nu})^{-2}(\hat{\sigma})^{-1}u \in L^2(0, T; H^2((\Omega)) \cap L^\infty(0, T; L^2(\Omega))) \hookrightarrow L^6(Q_T),$$

since we are working in an unidimensional case. At this point, we have fixed λ and μ such that

$$B_2 \subset L^2((\nu^*)^{-\frac{7}{2}}(\sigma^*)^{-1}(0, T); L^2(\Omega)) \times L^2((\nu^*)^{-\frac{7}{2}}(\sigma^*)^{-1}(0, T); H^{-2}(\Omega))^{2}$$

holds. Therefore, by using Hölder inequality, we get that

$$\begin{aligned} \left\| (\hat{\nu})^{-6}(\hat{\sigma})^{-3}u_1 u_2 u_3 \right\|_{L^2(Q_T)} &= \left\| ((\hat{\nu})^{-2}(\hat{\sigma})^{-1}u_1) ((\hat{\nu})^{-2}(\hat{\sigma})^{-1}u_2) ((\hat{\nu})^{-2}(\hat{\sigma})^{-1}u_3) \right\|_{L^2(Q_T)} \\ (3.17) \quad &\leq C \prod_{k=1}^3 \left\| (\hat{\nu})^{-2}(\hat{\sigma})^{-1}u_k \right\|_{L^6(Q_T)} \leq C \prod_{k=1}^3 \|(u_i, v_i, z_i)\|_{\mathcal{C}} \end{aligned}$$

and analogously, we have

$$\left\| (\hat{\nu})^{-6}(\hat{\sigma})^{-3}u_1 u_2 v_3 \right\|_{L^2(Q_T)} \leq C \prod_{k=1}^3 \|(u_i, v_i, z_i)\|_{\mathcal{C}},$$

showing the Claim 1.

Claim 2. $\mathcal{Y}'(0, 0, 0)$ is surjective.

First, note that $\mathcal{Y}(0, 0, 0) = (0, 0)$. By the other hand, observe that $\mathcal{Y}'(0, 0, 0) : B_1 \rightarrow B_2$ is given by

$$\mathcal{Y}'(0, 0, 0)(u, v, h) = (iu_t + u_{xx} - u_{xxxx} - 1_\omega h, iv_t + v_{xx} - v_{xxxx} - 1_{\mathcal{O}}u),$$

for $(u, v, h) \in B_1$. Invoking the null controllability result for linear system (1.6), that is, thanks to Theorem 3.3, $\mathcal{Y}'(0, 0, 0)$ is surjective, proving the Claim 2.

Finally, by taking $b_1 = (0, 0, 0)$, $b_2 = (0, 0)$ and using Theorem 3.4, there exists $\delta > 0$ such that if $\|(f^0, f^1)\|_{B_2} < \delta$, then we can find a control h such that the triple $(u, v, h) \in B_1$ satisfies $\mathcal{Y}(u, v, h) = (f^0, f^1)$. By a particular choice of $f^0 = f \in L^2((\hat{\nu})^{-6}(\hat{\sigma})^{-1}(0, T); L^2(\Omega))$ and $f^1 \equiv 0$, Theorem 1.2 is showed since a triple $(u, v, h) \in B_1$ satisfies $v(0) = 0$ in Ω and solves (1.5). \square

4. FURTHER COMMENTS AND OPEN ISSUES

To our knowledge, these results in this article are the first with respect to the existence of insensitizing controls for the fourth-order Schrödinger equation, in this way, we believe that this manuscript can open a series of questions, which are discussed now.

²Note that if necessary we could have taken λ and μ large enough in such a way that this inclusion is still satisfied.

4.1. Null condition of the initial data. In this point, we discuss the necessity to assume the null condition of the initial data in Theorem 1.2. In [11], the author proves that under some suitable conditions the existence of insensitizing controls may or may not hold, which indicates that this kind of problem cannot be solved for every initial data. In this way, we also have the same drawback in our result. To overcome this difficulty, we believe that the techniques used for Heat equation, due De Tereza [11], can be adapted for our case. Precisely, the idea consists in using the fundamental solution to construct an explicit solution where the observability inequality does not hold.

4.2. About the nonlinear terms. Note that if we change the cubic term $|u|^2u$ by a more general term $|u|^{p-2}u$, with $p \geq 3$, then one must prove a partial null controllability for the following system

$$\begin{cases} iu_t + u_{xx} - u_{xxxx} - \zeta|u|^{p-2}u = f + 1_\omega h, & \text{in } Q, \\ iv_t + v_{xx} - v_{xxxx} - \bar{\zeta}p|u|^{p-2}\bar{u}\bar{v} - (p+1)\bar{\zeta}|u|^{p-2}\bar{v} = 1_{\mathcal{O}}u, & \text{in } Q, \\ u(t, 0) = u(t, L) = v(t, 0) = v(t, L) = 0, & \text{on } t \in (0, T), \\ u_x(t, 0) = u_x(t, L) = v_x(t, 0) = v_x(t, L) = 0, & \text{on } t \in (0, T), \\ u(0, x) = u_0(x), v(T, x) = 0, & \text{in } \Omega. \end{cases}$$

If the structure of the problem is still the same and we only change the nonlinearity, the main difficult here is to obtain well-posedness results which gives enough regularity for the solutions to obtain the analogous Hölder estimate as in (3.17). In fact, to solve it one must have valid embedding from the state spaces into $L^{2p-2}(Q)$, for $p \geq 3$, which is possible since the following estimate holds

$$\| |u|^{p-2}u \|_{H^s(\Omega)} \leq C \|u\|_{H^s(\Omega)}^{p-1},$$

when $s \geq \frac{1}{2}$ and $p \geq 3$, see [25] for well-posedness of the general nonlinear problem.

Additionally, if we change to a general type nonlinearity g , we obtain the following optimal system

$$\begin{cases} iu_t + u_{xx} - u_{xxxx} + g(u) = f + 1_\omega h, & \text{in } Q, \\ iv_t + v_{xx} - v_{xxxx} + g'(u)v = 1_{\mathcal{O}}u, & \text{in } Q, \\ u(t, 0) = u(t, L) = v(t, 0) = v(t, L) = 0, & \text{on } t \in (0, T), \\ u_x(t, 0) = u_x(t, L) = v_x(t, 0) = v_x(t, L) = 0, & \text{on } t \in (0, T), \\ u(0, x) = u_0(x), v(T, x) = 0, & \text{in } \Omega. \end{cases}$$

It is expected that most of these problems have no solution, i.e., it is not possible to insensitize the functional unless we impose some conditions on g . To exemplify the comments above, some of these issues were already considered for the case of nonlinearities with superlinear growth at infinity. In [5], the authors dealing with a semilinear heat equation proved positive result of existence of insensitizing controls considering $g \in C^1$ a nonlinear function verifying $g'' \in L_{loc}^\infty(\mathbb{R})$, $g(0) = 0$ and

$$\lim_{|s| \rightarrow \infty} \frac{g'(s)}{\ln(1 + |s|)} = 0,$$

furthermore, the result is also valid for nonlinearities g of the form

$$|g(s)| = |p_1(s)| \ln^\alpha(1 + |p_2(s)|),$$

for all $|s| \geq s_0 > 0$, with $\alpha \in [0, 1)$ and p_i , $i = 1, 2$, are affine functions. Moreover, they proved negative results of existence considering a nonlinearity g verifying the conditions above, that is, taking g as

$$g(s) = \int_0^{|s|} \ln^\alpha(1 + |\sigma|^2) d\sigma, \quad \text{for all } s \in \mathbb{R},$$

but choosing $\alpha > 2$. Similar results are proved in [33] for a class of nonlinear Ginzburg-Landau equation.

Thus, in the case of the fourth-order nonlinear Schrödinger equation, this kind of situation, that is, introducing a function g with certain properties and to prove existence of insensitizing controls is still an open issue.

4.3. About the sentinel functional. One way to solve the problems of nonlinearity is to change the structure of the functional. Due to the lack of regularity of the characteristic function, if we change it to a more regular function then one can still prove result for more general nonlinearity $|u|^{p-2}u$, with $p \geq 3$, considering a functional of the form

$$J(\tau, h) = \frac{1}{2} \iint_{Q_T} \mathcal{R}(x) |u(x, t)|^2 dx dt,$$

where $\mathcal{R} \in C^\infty(\Omega)$ is a smooth function with $\text{Supp}(\mathcal{R}) \subset \mathcal{O}$.

We note that there exists an uncountable insensitizing control problems as we change the sentinel functional. In fact, by the equivalent formulation in a cascade system with double the equations of the original system, controllability problems with less control forces then equations are not fully understood in PDEs, so they can also be interesting from the control theory point of view. Some of the motivations for these problems arise from physical phenomena, thus typically we focus our attention in functionals which have “physical” meanings: If the functional is the local L^2 -norm of the the solution then we are looking for controls that locally preserve the energy (kinetic or potential, depending on the modeling) of the system, and if we change to a first derivative (or gradient in the N -dimensional case) the problem consists in finding controls that locally preserves the mean value of the energy.

In this perspective, let D be a derivative operator such as $Du = u_x$ or $Du = u_{xx}$. An interesting – and difficult – problem is to analyze the existence of insensitizing controls when the sentinel functional take the form

$$J(\tau, h) := \frac{1}{2} \int_{Q_T} |Du(x, t)|^2 dx dt.$$

In such, the optimal system become

$$\begin{cases} iu_t + u_{xx} - u_{xxxx} - \zeta |u|^2 u = f + 1_\omega h, & \text{in } Q_T, \\ iv_t + v_{xx} - v_{xxxx} - \bar{\zeta} \bar{u}^2 \bar{v} - 2\bar{\zeta} |u|^2 v = D(1_\mathcal{O} Du), & \text{in } Q_T, \\ u(t, 0) = u(t, L) = v(t, 0) = v(t, L) = 0, & \text{on } t \in (0, T), \\ u_x(t, 0) = u_x(t, L) = v_x(t, 0) = v_x(t, L) = 0, & \text{on } t \in (0, T), \\ u(0, x) = u_0(x), v(T, x) = 0, & \text{in } \Omega. \end{cases}$$

Again, it is not expected to obtain positive results of existence of insensitizing controls for every differential operator D in virtue of the lack of regularity provoked by coupling term $D(1_\mathcal{O} Du)$, since (again) the characteristic function is not regular. Despite that, Guerrero [19] dealt with a parabolic equation. The author proved a positive result of existence considering a functional depending on the gradient of the solution. Since the equation was linear, with constant coefficients, the argument consisted in considering a global Carleman estimate with different exponents, not for the equation, but for the equation satisfied by the Laplacian of the solutions to then recover information using the equation with the coupling. This is not the case when dealing with a nonlinear problem since deriving the equation would give us many other terms. In [21], the same author proved a similar result considering a linear Stokes equation with constant coefficients but with for the *curl* of the solution. Finally, we cite the work of the second author [22], where the authors proved positive results of insensitizing controls considering a functional depending on the gradient of the solution for the cubic nonlinear Ginzburg-Landau equation. The result arose by proving a new suitable Carleman estimate for the Ginzburg-Landau equation.

In this spirit, there are many alternatives to define the sentinel functional related with the insensitizing controls problems for 4NLS. Thus, we expect that these three works together with the results on this paper, open prospect to prove similar results considering a sentinel functional with the gradient of the solution. Moreover, since the Carleman estimate (2.3) has third order terms, maybe it is possible, at some point, to adapt the arguments to consider a functional with the Laplacian of the solution of the 4NLS, but clearly, to prove it is necessary new arguments of those that were applied here, at least proving a new Carleman estimate for the fourth-order Schrödinger equation, as was done in [18, Theorem 1.1] for the Cahn-Hilliard type equation. The readers are

invited to read the recent and interesting work by Imanuvilov and Yamamoto [20], which proves a Carleman estimate for a fourth-order parabolic equation in general dimensions.

4.4. N-dimensional case. Zheng and Zhou [34] studied the boundary controllability of the 4NLS in a bounded domain $\Omega \subset \mathbb{R}^n$. Using a L^2 -Neumann boundary control, the authors proved that the solution of 4NLS is exactly controllable in $H^{-2}(\Omega)$ using the Hilbert Uniqueness Method and multiplier techniques. In the sense of existence of insensitizing controls, we conjecture that the Carleman inequality shown here can be extended to the N -dimensional case. Thus, if we consider the sentinel functional as defined in (1.3), our result remains valid, for this case. However, the main issue here is when we consider a functional like the one mentioned in the Subsection 4.3 or other types of functional associated with the nonlinear problem. This type of problem looks interesting and still is open for the fourth-order nonlinear Schrödinger equation.

Acknowledgments. R. de A. Capistrano-Filho was supported by CNPq grant 307808/2021-1, CAPES grants 88881.311964/2018-01 and 88881.520205/2020-01, MATHAMSUD grant 21-MATH-03 and Propesqi (UFPE).

APPENDIX A. WELL-POSEDNESS

In this section we will show some results about the existence of solution for system

$$(A.1) \quad \begin{cases} iu_t + u_{xx} - u_{xxxx} = F^0, & \text{in } Q_T, \\ iv_t + v_{xx} - v_{xxxx} = F^1 + 1 \odot u, & \text{in } Q_T, \\ u(t, 0) = u(t, L) = u_x(t, 0) = u_x(t, L) = 0, & \text{on } t \in (0, T), \\ v(t, 0) = v(t, L) = v_x(t, 0) = v_x(t, L) = 0, & \text{on } t \in (0, T), \\ u(0, x) = u_0(x), v(T, x) = 0, & \text{in } \Omega, \end{cases}$$

for given u_0 and (F^0, F^1) . The proofs in here can be adapted to prove the existence of solutions for systems (1.6) and (2.4).

A.1. The linearized system. We first consider the simplest linear equation with null boundary conditions which is a linearized version of (1.2) around zero. More precisely, we consider the following

$$(A.2) \quad \begin{cases} iu_t + u_{xx} - u_{xxxx} = f, & \text{in } Q_T, \\ u(t, 0) = u(t, L) = u_x(t, 0) = u_x(t, L) = 0, & \text{on } (0, T), \\ u(0, x) = u_0(x), & \text{in } \Omega. \end{cases}$$

The first result is a consequence of the semigroup theory. Before presenting it, let us consider the differential operator $A : D(A) \subset L^2(\Omega) \rightarrow L^2(\Omega)$ given by

$$Au := iu_{xx} - iu_{xxxx},$$

with domain $D(A) = H^4(\Omega) \cap H_0^2(\Omega)$. Thus, the nonhomogeneous linear system (A.2) takes the form

$$(A.3) \quad \begin{cases} u_t(t) = Au(t) + if(t), & t \in [0, T], \\ u(0) = u_0. \end{cases}$$

The following proposition guarantees some properties for the operator A . Precisely, the result ensures the existence of regular solutions for the system (A.2).

Proposition A.1. *Let $f \in C^1([0, T]; L^2(\Omega))$ and $u_0 \in D(A)$, then (A.2) has a unique solution*

$$(A.4) \quad u \in C([0, T]; H^4(\Omega) \cap H_0^2(\Omega)) \cap C^1([0, T]; L^2(\Omega)).$$

Proof. Consider the linear operator defined by A . This allows us to rewrite (A.2) in the abstract form (A.3). We have that A is self-adjoint operator and A is m-dissipative. Indeed, first, is not difficult to see that

$$(Au, v)_{L^2(\Omega)} = (u, Av)_{L^2(\Omega)},$$

for all $u, v \in D(A)$. That is, A is symmetric. Additionally, $D(A^*) = D(A)$, so A is self-adjoint. Finally, we have

$$(Au, u)_{L^2(\Omega)} = \operatorname{Re} \left(i \int_{\Omega} (u_{xx} - u_{xxxx}) \bar{u} dx \right) = \operatorname{Re} \left(i \int_{\Omega} -(|u_x|^2 + |u_{xx}|^2) dx \right) = 0,$$

for any $u \in D(A)$, and then A is dissipative. Therefore, A is a m-dissipative operator (e.g. [9, Corollary 2.4.8]) and by the Hille–Yosida–Phillips theorem (e.g. [9, Theorem 3.4.4]) we obtain that A is a generator of a contraction semigroup in $L^2(\Omega)$. Thus, if $u_0 \in D(A)$ and $f \in C^1([0, T]; L^2(\Omega))$, then equation (A.2) has solutions u with the regularity (A.4). (e.g. [9, Proposition 4.1.6]). \square

A.2. The coupled linearized system. We are now concerned with the existence of solutions for the coupled linearized system. More precisely, we will prove the well-posedness results to the system (A.1). First, consider the linear unbounded operator A defined in the previous subsection and

$$\begin{cases} A_1 u = -iu_{xx} + iu_{xxxx} \in H^{-2}(\Omega), \\ D(A_1) = H_0^2(\Omega). \end{cases}$$

Both operators are m-dissipative with dense domains; therefore, they generate the C_0 semigroups of contractions \mathcal{S}_0 and \mathcal{S}_1 , respectively. Now, consider the following spaces:

$$Y_0 = C([0, T]; D(A)) \cap C^1([0, T]; L^2(\Omega))$$

and

$$Y_1 = C([0, T]; D(A_1)) \cap C^1([0, T]; H^{-2}(\Omega)).$$

The next result is dedicated to prove the existence of *regular solutions* for (A.1)

Theorem A.2 (Regular solutions). *Assume that $u_0 \in D(A)$,*

$$F^0 \in C([0, T], H^4(\Omega) \cap H_0^2(\Omega)) \cap W^{1,1}(0, T; H^4(\Omega) \cap H_0^2(\Omega))$$

and

$$F^1 \in C([0, T], H^{-2}(\Omega)) \cap W^{1,1}(0, T; H^{-2}(\Omega)).$$

Then, problem (A.1) has a unique regular solution in the sense that

$$\begin{cases} (u, v) \in Y_0 \times Y_1, \\ iu_t + u_{xx} - u_{xxxx} = F^0, \\ iv_t + v_{xx} - v_{xxxx} = F^1 + 1_{\mathcal{O}}u, \\ u|_{t=0} = u_0, v|_{t=T} = 0. \end{cases}$$

Proof. Note that, thanks to [9, Proposition 4.1.6], we get that the mild solution

$$u(t) = \mathcal{S}_0(t)u_0 + \int_0^t \mathcal{S}_0(t-s)F^0(s)ds \in Y_0$$

verifies

$$(A.5) \quad \begin{cases} iu_t + u_{xx} - u_{xxxx} = F^0, \\ u|_{t=0} = u_0. \end{cases}$$

Now, it is not difficult to see that

$$\int_0^T \|(u1_{\mathcal{O}})(s)\|_{H^{-2}(\Omega)}^2 ds \leq \sup_{\zeta \in H_0^2(\Omega), \|\zeta\|_{H_0^2(\Omega)}=1} \int_0^T \int_{\mathcal{O}} |u| \cdot \zeta dx dt \leq \iint_{Q_T} |u|^2 dx dt,$$

and hence $u1_{\mathcal{O}} \in C([0, T], H^{-2}(\Omega)) \cap W^{1,1}(0, T; H^{-2}(\Omega))$. Then, applying again [9, Proposition 4.1.6], and we get that the mild solution

$$v(t) = \int_t^T \mathcal{S}_1(s-t)(F^1 + 1_{\mathcal{O}}u)(s)ds \in Y_1$$

satisfies

$$(A.6) \quad \begin{cases} iv_t + v_{xx} - v_{xxxx} = F^1 + 1_{\mathcal{O}}u, \\ v|_{t=T} = 0. \end{cases}$$

Thus, Theorem A.2 is achieved putting together (u, v) satisfying (A.5) and (A.6). \square

A.3. Transposition solutions. In what follows, we will talk about transposition solutions that are of particular interest for the purposes of this paper.

Definition 2. Let $u_0 \in L^2(\Omega)$ and $(F^0, F^1) \in [L^2(0, T; H^{-2}(\Omega))]^2$. We say that a pair

$$(u, v) \in L^2(0, T; H_0^2(\Omega)) \times L^2(Q_T)$$

is a solution in the transposition sense of (A.1), if it satisfies

$$(A.7) \quad \begin{aligned} \int_0^T \langle g^0, u \rangle_{H^{-2}H_0^2} dt &= \operatorname{Re} \int_{\Omega} \varphi(0) \overline{u_0} dx + \int_0^T \langle F^0, \varphi \rangle_{H^{-2}H_0^2} dt \\ \operatorname{Re} \iint_{Q_T} g^1 \bar{v} dx dt &= \operatorname{Re} \int_0^T \int_{\mathcal{O}} u \cdot \bar{\psi} dx dt + \int_0^T \langle F^1, \psi \rangle_{H^{-2}H_0^2} dt \end{aligned}$$

for every $(g^0, g^1) \in L^2(0, T; H^{-2}(\Omega)) \times L^2(Q_T)$, where $\langle \cdot, \cdot \rangle$ denotes the duality between $H^{-2}(\Omega)$ and $H_0^2(\Omega)$, and (φ, ψ) is the solution of

$$(A.8) \quad \begin{cases} i\varphi_t + \varphi_{xx} - \varphi_{xxxx} = g^0, & \text{in } Q_T, \\ i\psi_t + \psi_{xx} - \psi_{xxxx} = g^1, & \text{in } Q_T, \\ \varphi(t, 0) = \varphi(t, L) = \varphi_x(t, 0) = \varphi_x(t, L) = 0, & \text{on } t \in (0, T), \\ \psi(t, 0) = \psi(t, L) = \psi_x(t, 0) = \psi_x(t, L) = 0, & \text{on } t \in (0, T), \\ \varphi(T, x) = 0, \psi(0, x) = 0, & \text{in } \Omega. \end{cases}$$

We have the following result about the existence and uniqueness of transposition solutions.

Theorem A.3. For $u_0 \in L^2(\Omega)$ and $(F^0, F^1) \in [L^2(0, T; H^{-2}(\Omega))]^2$, there exists a unique $(u, v) \in [L^2(0, T; H_0^2(\Omega))]^2$ satisfying (A.7) for every $(g^0, g^1) \in L^2(0, T; H^{-2}(\Omega)) \times L^2(Q_T)$, where (φ, ψ) is solution of (A.8).

Proof. Let $\Psi_1 : L^2(0, T; H_0^2(\Omega)) \rightarrow \mathbb{R}$ the operator defined by

$$\Psi_1(h^0) = \operatorname{Re} \int_{\Omega} \varphi(0) \overline{u_0} dx + \int_0^T \langle F^0, \varphi \rangle_{H^{-2}H_0^2} dt,$$

where φ satisfies the first equation of (A.8) for $g^0 := h_{xxxx}^0 \in L^2(0, T; H^{-2}(\Omega))$. From the energy estimates, it is easy to see the continuity of Ψ_1 . Then, thanks to the Lax-Milgran theorem, there exists $u \in L^2(0, T; H_0^2(\Omega))$ such that

$$\int_0^T \langle g^0, u \rangle_{H^{-2}H_0^2} dt = \operatorname{Re} \iint_{Q_T} h_{xx}^0 \bar{u}_{xx} dx dt = \Psi_1(h^0),$$

for every $g^0 \in H^{-2}(\Omega)$, with $g^0 = h_{xxxx}^0$. Analogously, we have the existence of $v \in L^2(Q_T)$ satisfying the second equation of (A.7), since the linear form

$$\Psi_1(g^1) = \operatorname{Re} \int_0^T \int_{\mathcal{O}} u \cdot \bar{\psi} dx dt + \int_0^T \langle F^1, \psi \rangle_{H^{-2}H_0^2} dt$$

is continuous in $L^2(Q_T)$.

Claim: We have that v belongs to $L^2(0, T; H_0^2(\Omega))$.

Indeed, first, we take sequences of regular data such that $u_0^n \rightarrow u_0$ in $L^2(\Omega)$ and $(F_n^0, F_n^1) \rightarrow (F^0, F^1)$ in $L^2(0, T; H^{-2}(\Omega)) \times L^2(Q_T)$. We show that the regular solutions (u_n, v_n) for (A.1) (whose existence is given in Theorem A.2) with initial data u_0^n and (F_n^0, F_n^1) on the right-hand

side, are also a solution in the transposition sense; moreover, it is bounded in $[L^2(0, T; H_0^2(\Omega))]^2$. Hence, in the limit, we obtain that $(u, v) \in [L^2(0, T; H_0^2(\Omega))]^2$.

Finally, for uniqueness, suppose that (\hat{u}, \hat{v}) is another solution of (A.1). Thus,

$$\operatorname{Re} \int_0^T \langle g^0, u - \hat{u} \rangle_{H^{-2}H_0^2} dt = 0 \quad \text{and} \quad \operatorname{Re} \iint_{QT} g^1(v - \hat{v}) dx dt = \operatorname{Re} \int_0^T \int_{\mathcal{O}} (u - \hat{u}) \cdot \bar{\psi} dx dt,$$

for all $g^0 \in L^2(0, T; H^{-2}(\Omega))$ and $g^1 \in L^2(Q_T)$. Hence, $y = \hat{y}$ and $z = \hat{z}$. \square

REFERENCES

- [1] F. ALABAU-BOUSSOIRA, *Insensitizing exact controls for the scalar wave equation and exact controllability of coupled cascade systems of PDE's by a single control*. Math. Control Signals Syst. 26, 1–46 (2014).
- [2] V. M. ALEKSEEV, V. M. TIKHOMIROV AND S. V. FOMIN, *Optimal Control*. Translated from the Russian by V. M. Volosov, Contemporary Soviet Mathematics Consultants Bureau, New York (1987).
- [3] M. BEN-ARTZI, H. KOCH AND J.-C. SAUT, *Dispersion estimates for fourth order Schrödinger equations*. C. R. Acad. Sci. Paris Sér. I Math 330:87–92 (2000).
- [4] O. BODART AND C. FABRE, *Controls Insensitizing the Norm of the Solution of a Semilinear Heat-Equation*. Journal of mathematical analysis and applications, 195, 3:658–683 (1995).
- [5] O. BODART, M. GONZÁLEZ-BURGOS AND R. PÉREZ-GARCÍA, ROSARIO, *Existence of insensitizing controls for a semilinear heat equation with a superlinear nonlinearity*. Commun. Partial Differ. Equ. 29(7–9), 1017–1050 (2004).
- [6] O. BODART, M. GONZÁLEZ-BURGOS AND R. PÉREZ-GARCÍA, ROSARIO, *Insensitizing controls for a heat equation with a nonlinear term involving the state and the gradient*. Nonlinear Analysis: Theory, Methods & Applications. (5-6):687–711 (2004).
- [7] B. CALSAVARA, N. CARREÑO AND E. EDUARDO, *Insensitizing controls for a phase field system*. Nonlinear Analysis: Theory, Methods & Applications. 143:120–137 (2016).
- [8] N. CARREÑO, *Insensitizing controls for the Boussinesq system with no control on the temperature equation*. Advances in Differential Equations 22 (3/4):235–258 (2017).
- [9] T. CAZENAVE AND A. HARAUX, *An Introduction to Semilinear Evolution Equations*, in: Oxford Science Publications, Oxford University Press, 1998.
- [10] L. DE TERESA, *Controls insensitizing the norm of the solution of a semilinear heat equation in unbounded domains*. Esaim-control Optimisation and Calculus of Variations. 2:125–149 (1997).
- [11] L. DE TERESA, *Insensitizing controls for a semilinear heat equation: Semilinear heat equation*. Communications in Partial Differential Equations, 25 (1-2): 39–72 (2000).
- [12] L. DE TERESA, M. LOPEZ-GARCIA AND A. MERCADO, *Null controllability of a cascade system of Schrödinger equations*. Electronic Journal of Differential Equations. 74:1–12 (2016).
- [13] L. DE TEREZA, S. MICU AND J. ORTEGA, *An example of ϵ -insensitizing controls for the heat equation with no intersecting observation and control regions*. Appl. Math. Lett. 17:927–932 (2004).
- [14] L. DE TERESA AND E. ZUAZUA, *Identification of the class of initial data for the insensitizing control of the heat equation*. Commun. Pure Appl. Anal 8 (1):457–471 (2009).
- [15] S. DOLECKI AND D. L. RUSSELL, *A general theory of observation and control*. SIAM J. Control Opt. 15:185–220 (1977).
- [16] E. FERNANDEZ-CARA, G. GARCIA AND A. OSSER, *Insensitizing controls for a large-scale ocean circulation model*. Comptes Rendus Mathématique. 337:265–270 (2003).
- [17] P. GAO, *Insensitizing controls for the Cahn-Hilliard type equation*. Electronic Journal of Qualitative Theory of Differential Equations. 35:1–22 (2018).
- [18] P. GAO, *A new global Carleman estimate for the Cahn-Hilliard type equation and its applications*. Journal of Differential Equations. 260(1):427–444 (2016).
- [19] S. GUERRERO, *Null controllability of some systems of two parabolic equations with one control force*. SIAM Journal on Control and Optimization. 46 (2):379–394 (2007).
- [20] O. YU. IMANUVILOV AND M. YAMAMOTO, *Inverse source problem and the continuation for a fourth-order parabolic equation in general dimensions*. arXiv:2009.11592v1 [math.AP].
- [21] S. GUERRERO, *Controllability of systems of Stokes equations with one control force: existence of insensitizing controls*, Annales de l'Institut Henri Poincaré (C) Non Linear Analysis. 24 (6):1029–1054 (2007).
- [22] M. C. SANTOS AND T. Y. TANAKA, *An Insensitizing Control Problem for the Ginzburg-Landau Equation*. J Optim Theory Appl. 183:440–470 (2019).
- [23] V. I. KARPMAN, *Stabilization of soliton instabilities by higher-order dispersion: fourth order nonlinear Schrödinger-type equations*. Phys. Rev. E. 53:1336–1339 (1996).
- [24] V. I. KARPMAN AND A. G. SHAGALOV, *Stability of soliton described by nonlinear Schrödinger type equations with higher-order dispersion*. Physica D. 144:194–210 (2000).

- [25] J. LI AND C. ZHENG, *Non-homogeneous initial boundary value problems for the biharmonic Schrödinger equation on an interval*. arXiv:2003.09337v3 [math.FA].
- [26] J.-L. LIONS, *Exact controllability, stabilization and perturbations for distributed systems*. SIAM Rev. 30:1–68 (1988).
- [27] J.-L. LIONS, *Quelques notions dans l'analyse et le contrôle de systèmes à données incomplètes*. Proceedings of the XIth Congress on Differential Equations and Applications/First Congress on Applied Mathematics (Spanish)(Málaga, 1989). 43–54 (1990).
- [28] J.-L. LIONS, *Sentinelles pour les systèmes distribués à données incomplètes*. vol. 21. Masson, Paris (1992).
- [29] P. LISSY, Y. PRIVAT AND Y. SIMPONE, *Insensitizing control for linear and semi-linear heat equations with partially unknown domain*. ESAIM: Control, Optimisation and Calculus of Variations. 25:50 (2019).
- [30] B. PAUSADER, *The cubic fourth-order Schrödinger equation*. J. Funct. Anal. 256:2473–2517 (2009).
- [31] R. K. RAJAGOUNDER AND C. K. TO, *Insensitizing control of KDV-Burgers Equations*. Journal of Electrical Systems and Information Technology. 5:2 192–207 (2018).
- [32] M. ZHANG, J. YIN AND H. GAO, *Insensitizing controls for the parabolic equations with dynamic boundary conditions*. Journal of Mathematical Analysis and Applications. 475:1 861–873 (2019).
- [33] M. ZHANG AND X. LIU, *Insensitizing controls for a class of nonlinear Ginzburg-Landau equation*. Sci. China Math. 57(12), 2635–2648 (2014).
- [34] C. ZHENG, *Inverse problems for the fourth order Schrödinger equation on a finite domain*, Mathematical Control & Related Fields, 5(1): 177–189 (2015).

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE PERNAMBUCO (UFPE), 50740-545, RECIFE (PE), BRAZIL.

Email address: roberto.capistranofilho@ufpe.br

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL RURAL DE PERNAMBUCO (UFRPE), 52171-900, RECIFE (PE), BRAZIL.

Email address: thiago.tanaka@ufrpe.br